



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(1) International Patent Classification 6 : C07D 311/58, A61K 31/495, 31/535		AI	(11) International Publication Number: WO 97/4212 (43) International Publication Date: 25 March 1999 (25.03.99)
(21) International Application Number: PCT/SE98/01603 (22) International Filing Date: 9 September 1998 (09.09.98) (30) Priority Data: 970377-3 18 September 1997 (18.09.97) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG [SE/SE]: S-151 85 Södertälje (SE). (72) Inventors, and (73) Invention/Applicants (for US only): BERO, Stefan [SE/SE]; Astra Actus AB, S-151 85 Södertälje [SE/AT/PT/JP/US/SE]; Astra Actus AB, S-151 85 Södertälje [SE/AT/PT/JP/US/SE]; Astra Actus AB, S-151 85 Södertälje [SE]; Astra Actus AB, S-151 85 Södertälje [SE]; Astra Actus AB, S-151 85 Södertälje [SE]; Astra Actus AB, S-151 85 Södertälje [SE]. (74) Agent: ASTRA AKTIEBOLAG, Patent Dept., S-151 85 Södertälje (SE).		Published with international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: SUBSTITUTED CHROMAN DERIVATIVES			
(37) Abstract <p>The present invention relates to new piperidin- or piperazin-antibiotic dihydro-2H-1-benzopyran derivatives compounds having formula (I) wherein X is N or CH; Y is NR²CH₂, CH₂NR², NR²CO, CONR², NR²SO, or NR²CONR², wherein R₁ is H or C₁-C₄ alkyl; R₂ is H or C₁-C₄ alkyl or C₃-C₆ cycloalkyl; R₃ is C₁-C₄ alkyl; C₃-C₆ cycloalkyl or (CH₂)_n-aryl, wherein aryl is phenyl or a heteroaromatic ring containing one or two heteroatoms selected from N, O and S and which may be mono- or di-substituted with R₄ and/or R₅, and n is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.</p>			

C1CCN(C1)c2ccc3c(c2)ccc(O3)C4CC()CC4

C

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT

FOR THE PURPOSES OF INFORMATION ONLY

[illegible]

SUBSTITUTED CHROMAN DERIVATIVES

Field of the Invention

The present invention relates to new piperidyl- or piperazinyl-substituted dihydro-2H-1-benzopyran derivatives as (*R*)- enantiomers, (*S*)-enantiomers or racemates in the form of free base or pharmaceutically acceptable salts or solvates thereof, a process for their preparation, pharmaceutical compositions containing said therapeutically active compounds and to the use of said active compounds in therapy.

An object of the invention is to provide compounds for therapeutic use, especially compounds having a selective effect at a subgroup of 5-hydroxytryptamine receptors, designated the h5-HT_{1B}-receptor (previously called the 5-HT_{1D_B}-receptor) in mammals including man.

It is also an object of the invention to provide compounds with a therapeutic effect after oral administration.

Background of the Invention

Various central nervous system disorders such as depression, anxiety, etc. appear to involve the disturbance of the neurotransmitters noradrenaline (NA) and 5-hydroxytryptamine (5-HT), the latter also known as serotonin. The drugs most frequently used in the treatment of depression are believed to act by improving the neurotransmission of either or both of these physiological agonists. It appears that the enhancement of 5-HT neurotransmission primarily affects the depressed mood and anxiety, whereas the enhancement of noradrenaline neurotransmission affects the retardation symptoms occurring in depressed patients. The invention concerns compounds which have an effect on 5-HT neurotransmission.

Serotonin, or 5-HT, activity is believed to be involved in many different types of psychiatric disorders. For instance it is believed that an increase in 5-HT activity is

associated with anxiety, while a decrease in 5-HT release has been associated with depression. Serotonin has in addition been implicated in such diverse conditions as eating disorders, gastrointestinal disorders, cardiovascular regulation disorders and sexual disturbances.

The 5-HT Receptors

The various effects of 5-HT may be related to the fact that serotonergic neurons stimulate the secretion of several hormones, e.g. cortisol, prolactin, β -endorphin, vasopressin and others. The secretion of each of these other hormones appears to be regulated on a specific basis by several different 5-HT (serotonin) receptor subtypes. With the aid of molecular biology techniques, to date these receptors have been classified as 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆ and 5-HT₇ with the 5-HT₁ receptor further divided into the 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E} and 5-HT_{1F} subtypes. Each receptor subtype is involved in a different serotonin function and has different properties.

Regulation of the 5-HT transmission

The release of 5-HT is feedback-regulated by two different subtypes of 5-HT receptors. Inhibitory 5-HT_{1A} autoreceptors are located on the cell bodies in the raphe nuclei which upon stimulation by 5-HT decrease the impulse propagation in the 5-HT neurons and thereby reducing the 5-HT released at the nerve terminals. Another subtype of inhibitory 5-HT receptors is located on the 5-HT nerve terminals, the h5-HT_{1B} receptors (in rodents the r5-HT_{1B} receptors) which regulate the synaptic concentration of 5-HT by controlling the amount of 5-HT that is released. An antagonist of these terminal autoreceptors thus increases the amount of 5-HT released by nerve impulses which has been shown in both *in vitro* and *in vivo* experiments.

The use of an antagonist of the terminal h5-HT_{1B} autoreceptor will accordingly increase the synaptic 5-HT concentration and enhance the transmission in the 5-HT system. It would thus produce an antidepressant effect making it useful as a medication for depression.


Other localizations of h5-HT_{1B} receptor subtype also exist. A large part of these postsynaptic receptors appear to be located on nerve terminals of other neuronal systems (so called heteroreceptors). Since the h5-HT_{1B} receptor mediates inhibitory responses an antagonist of this receptor subtype might also increase the release of other neurotransmitters than 5-HT.

Compounds having h5-HT_{1B} activity may according to well known and recognised pharmacological tests be divided into full agonists, partial agonists and antagonists.

Disclosure of the Invention

The object of the present invention is to provide compounds having a selective effect at the h5-HT_{1B} receptor, preferably antagonistic properties, as well as having a good bioavailability. The effect on the other receptors chosen from, for example, the 5-HT_{1A}, 5-HT_{2A}, D₁, D_{2A}, D₃, α_1 and α_2 receptor has been investigated.

Accordingly, the present invention provides compounds of the formula I



(I)

wherein

X is N or CH;

Y is NR₂CH₂, CH₂NR₂, NR₂CO, CO-NR₂, NR₂SO₂ or NR₂CONR₂

wherein R₂ is H or C₁-C₆ alkyl;

R₁ is H, C₁-C₆ alkyl or C₃-C₆ cycloalkyl;

R₃ is C₁-C₆ alkyl, C₃-C₆ cycloalkyl or (CH₂)_n-aryl,

wherein aryl is phenyl or a heteroaromatic ring containing one or two heteroatoms selected from N, O and S and which may be mono- or di-substituted with R₄ and/or R₅:

wherein R₄ is H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, halogen, CN, CF₃, OH, C₁-C₆ alkoxy, NR₆R₇, OCF₃, SO₂CH₃, SO₂CF₃, SO₂NR₆R₇, phenyl, phenyl-C₁-C₆ alkyl, phenoxy, C₁-C₆ alkylphenyl, an optionally substituted heterocyclic ring containing one or two heteroatoms selected from N, O, S, SO and SO₂ wherein the substituent(s) is(are) selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl, phenyl-C₁-C₆ alkyl, (CH₂)_mOR₉ wherein m is 2-6 and R₉ is H, C₁-C₆ alkyl, C₃-C₆ cycloalkyl or phenyl-C₁-C₆ alkyl, and COR₈, an optionally substituted heteroaromatic ring containing one or two heteroatoms selected from N, O and S wherein the substituent(s) is(are) selected from C₁-C₆ alkyl, C₃-C₆ cycloalkyl and phenyl-C₁-C₆ alkyl, or COR₈:

wherein R₆ is H, C₁-C₆ alkyl or C₃-C₆ cycloalkyl;

R₇ is H, C₁-C₆ alkyl or C₃-C₆ cycloalkyl; and

R₈ is C₁-C₆ alkyl, C₃-C₆ cycloalkyl, CF₃, NR₆R₇, phenyl, a

heteroaromatic ring containing one or two heteroatoms selected from

N, O and S or a heterocyclic ring containing one or two heteroatoms

selected from N, O, S, SO and SO₂;

R₅ is H, OH, CF₃, OCF₃, halogen, C₁-C₆ alkyl or C₁-C₆ alkoxy;

and n is 0-4;

as (R)-enantiomers, (S)-enantiomers or a racemate in the form of a free base or a pharmaceutically acceptable salt or solvate thereof which possess a high selective effect at the h5-HT_{1B} receptor and also show sufficient bioavailability after oral administration.

In the present context C₁-C₆ alkyl may be straight or branched. C₁-C₆ alkyl may be methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, t-pentyl, neo-pentyl, n-hexyl or i-hexyl.

5 In the present context C₁-C₆ alkoxy may be straight or branched. C₁-C₆ alkoxy may be methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, i-butoxy, s-butoxy, t-butoxy, n-pentyloxy, i-pentyloxy, t-pentyloxy, neo-pentyloxy, n-hexyloxy or i-hexyloxy.

10 In the present context C₃-C₆ cycloalkyl may be cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, preferably cyclohexyl.

In the present context halogen may be fluoro, chloro, bromo or iodo.

15 In the present context the heteroaromatic ring containing one or two heteroatoms selected from N, O and S preferably is a 5- or 6-membered heteroaromatic ring and may be furyl, imidazolyl, isoxazolyl, isobiazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, thiazolyl or thienyl. The heteroaromatic ring can be either substituted or unsubstituted.

20 In the present context the heterocyclic ring containing one or two heteroatoms selected from N, O, S, SO and SO₂ may optionally contain a carbonyl function and is preferably a 5-, 6- or 7-membered heterocyclic ring and may be imidazolidinyl, imidazolyl, morpholinyl, piperazinyl, piperidyl, piperidinyl, pyrazolidinyl, pyrazolyl, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, thiomorpholinyl, preferably piperidino, 1-piperazinyl, morpholino, thiomorpholino and 4-piperidino-1-yl.

25 A preferred embodiment of the invention relates to compounds of formula I wherein Y is NHCO or CONH, i.e. amides. Of these compounds, the compounds wherein R₃ is unsubstituted phenyl, or mono- or di- substituted phenyl, and especially ortho-, meta- or para- substituted phenyl, and particularly these wherein the substituent R₄ is phenyl,

phenyl-C₁-C₆ alkyl, cyclohexyl, piperidino, 1-piperazinyl, morpholino, CF₃, 4-piperidino-1-yl, n-butoxy or COR₈ wherein R₈ is phenyl, cyclohexyl, 4-piperidino-1-yl, 1-piperazinyl, morpholino, CF₃, piperidino or NR₆R₇, are preferred.

5 Examples of combinations of substituents are:

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

10 X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is hydroxyethyl-piperazinyl, R₅ is H;

15 X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl;

20 X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is morpholino, R₅ is H;

25 X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is morpholino, R₅ is H;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is morpholino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is morpholino, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is piperidino, R₅ is H;

5 X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is hydroxyethyl-piperazinyl, R₅ is H;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is morpholino, R₅ is H;

10 X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is benzyloxyethyl-piperazinyl, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl;

15 X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is piperidino, R₅ is H;

20 X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

25 X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is morpholino, R₅ is H;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl;

30 X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is morpholino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is benzyloxyethyl-piperazinyl, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is COR₈, R₅ is morpholino;

5 X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is morpholino, R₅ is H;

10 X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is piperidino, R₅ is H;

15 X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is COR₈, R₅ is NR₈R₇, R₈R₇CH₃, C₂H₅ or C₃H₇;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

20 X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is morpholino, R₅ is H;

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl; -

X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

25 X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is morpholino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is COR₈, R₅ is morpholino;

30 X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is morpholino, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperaziny, R₅ is H;

5 X is CH, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is piperidino, R₅ is H;

10 X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl, R₄ is phenyl, phenylmethyl or phenylethyl, R₅ is H;

X is CH, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is piperidino, R₅ is H;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is (CH₂)₂-phenyl, R₄ is morpholino, R₅ is H;

15 X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is piperaziny, R₅ is H;

X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl, R₄ is COR₈, R₈ is cyclohexyl;

20 X is N, Y is CONR₂, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is phenyl;

X is N, Y is NR₂CO, R₁ is H, CH₃, C₂H₅ or C₃H₇, R₂ is H, R₃ is CH₂-phenyl.

Preferred compounds are:

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-

morpholinobenzamide

25 (*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-piperidinobenzamide

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-butoxybenzamide

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-

trifluoromethylbenzamide

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-*N,N*-diethylaminobenzamide

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-

trifluoromethoxybenzamide

5 (*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-piperidin-1-yl)benzamide

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-tetrahydro-1,4-diazepin-5-on-1-yl)benzamide, and

(*S*)-*N*-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran-3-yl]-4-(4-

10 benzylpiperazin-1-yl)benzamide.

The compounds of the present invention are in the form of the racemate or the (*R*)- or (*S*)-enantiomer in the form of a free base or a pharmaceutically acceptable salt or solvate thereof. Compounds in the form of the (*S*)-enantiomer are preferred ones.

15 Both organic and inorganic acids can be employed to form non-toxic pharmaceutically acceptable acid addition salts of the compounds of this invention. Illustrative acids are sulfuric, nitric, phosphoric, oxalic, hydrochloric, formic, hydrobromic, citric, acetic, lactic, tartaric, dibenzoyltartaric, diacetyltartaric, palmitic, ethanedisulfonic, sulfamic, succinic, propionic, glycolic, malic, gluconic, pyruvic, phenylacetic, 4-aminobenzoic, anthranilic, salicylic, 4-aminosalicylic, 4-hydroxybenzoic, 3,4-dihydroxybenzoic, 3,5-

20 dihydroxybenzoic, 3-hydroxy-2-naphthoic, nicotinic, methanesulfonic, ethanesulfonic, hydroxyethanesulfonic, benzenesulfonic, p-toluenesulfonic, sulfamic, naphthalenesulfonic, ascorbic, cyclohexylsulfamic, fumaric, maleic and benzoic acids. These salts are readily prepared by methods known in the art.

The preferred solvates of the compounds of this invention are the hydrates.

11

Pharmaceutical Formulations

In a second aspect the present invention provides a pharmaceutical formulation comprising as active ingredient a therapeutically effective amount of the compound of formula I as an enantiomer or a racemate in the form of a free base or a pharmaceutically acceptable salt or solvate thereof, optionally in association with diluents, excipients or inert carriers.

According to the present invention the compound of the invention will normally be administered orally, rectally or by injection, in the form of pharmaceutical formulations comprising the active ingredient either as a free base or a pharmaceutically acceptable non-toxic acid addition salt, e.g. the hydrochloride, hydrobromide, lactate, acetate, phosphate, sulfate, sulfamate, citrate, tartrate, oxalate and the like in a pharmaceutically acceptable dosage form. The dosage form may be a solid, semisolid or liquid preparation. Usually the active substance will constitute between 0.1 and 99% by weight of the preparation, more specifically between 0.5 and 20% by weight for preparations intended for injection and between 0.2 and 50% by weight for preparations suitable for oral administration.

To produce pharmaceutical formulations containing the compound of the invention in the form of dosage units for oral application, the selected compound may be mixed with a solid excipient, e.g. lactose, saccharose, sorbitol, mannitol, starches such as potato starch, corn starch or amylopectin, cellulose derivatives, a binder such as gelatine or polyvinylpyrrolidone, and a lubricant such as magnesium stearate, calcium stearate, polyethylene glycol, waxes, paraffin, and the like, and then compressed into tablets. If coated tablets are required, the cores, prepared as described above, may be coated with a concentrated sugar solution which may contain, e.g. gum arabic, gelatine, talcum, titanium dioxide, and the like. Alternatively, the tablet can be coated with a polymer known to the person skilled in the art, dissolved in a readily volatile organic solvent or mixture of organic solvents. Dyes/ultraviolet may be added to these coatings in order to readily distinguish between tablets containing different active substances or different amounts of the active compound.

12

For the preparation of soft gelatine capsules, the active substance may be admixed with e.g. a vegetable oil or polyethylene glycol. Hard gelatine capsules may contain granules of the active substance using either the above mentioned excipients for tablets, e.g. lactose, saccharose, sorbitol, mannitol, starches (e.g. potato starch, corn starch or amylopectin), cellulose derivatives or gelatine. Also liquids or semisolids of the drug can be filled into hard gelatine capsules.

Dosage units for rectal application can be solutions or suspensions or can be prepared in the form of suppositories comprising the active substance in a mixture with a neutral fatty base, or gelatine rectal capsules comprising the active substance in admixture with vegetable oil or paraffin oil. Liquid preparations for oral application may be in the form of syrups or suspensions, for example solutions containing from about 0.1% to about 20% by weight of the active substance herein described, the balance being sugar and mixture of ethanol, water, glycerol and propylene glycol. Optionally such liquid preparations may contain colouring agents, flavouring agents, saccharine and carboxymethyl-cellulose as a thickening agent or other excipients known to the person skilled in the art.

Solutions for parenteral applications by injection can be prepared in an aqueous solution of a water-soluble pharmaceutically acceptable salt of the active substance, preferably in a concentration of from about 0.1% to about 10% by weight. These solutions may also contain stabilizing agents and/or buffering agents and may conveniently be provided in various dosage unit ampoules.

Suitable daily doses of the compound of the invention in therapeutic treatment of humans are about 0.01-100 mg/kg bodyweight at peroral administration and 0.001-100 mg/kg bodyweight at parenteral administration.

The compound of the invention may be used in a combination with a 5-HT₂ receptor inhibitor, such as fluoxetine, paroxetine, citalopram, clomipramine, sertraline, alprazolam or fluvoxamin, preferably paroxetine or citalopram. Another possible combination is to use

the compound of the invention together with a monoamine oxidase inhibitor, such as moclobemide, tranylcypromine, brofaromide or phenelzine, preferably moclobemide or phenelzine. Still another possible combination is the compound of the invention together with a 5-HT_{1A} antagonist, such as the compounds disclosed in WO 96/33710, preferably (*R*)-5-carbamoyl-3-(*N,N*-dicyclobutylamino)-8-fluoro-3,4-dihydro-2*H*-1-benzopyran.

Medical and Pharmaceutical Use

In a further aspect the present invention provides the use of the compounds of formula I in therapy as a 5-HT_{1B} antagonist, partial agonist or full agonist, preferably as an antagonist and the use in the treatment of 5-hydroxytryptamine mediated disorders. Examples of such disorders are disorders in the CNS such as mood disorders (depression, major depressive episodes, dysthymia, seasonal affective disorder, depressive phases of bipolar disorder), anxiety disorders (obsessive compulsive disorder, panic disorder with/without agoraphobia, social phobia, specific phobia, generalized anxiety disorder, posttraumatic stress disorder), personality disorders (disorders of impulse control, trichotellomania), obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders (age associated memory impairment, presenile and senile dementia), pathological aggression, schizophrenia, endocrine disorders (e.g. hyperprolactinaemia), stroke, dyskinesia, Parkinson's disease, thermoregulation, pain and hypertension. Other examples of 5-hydroxytryptamine mediated disorders are urinary incontinence, vasospasm and growth control of tumors (e.g. lung carcinoma).

Methods of Preparation


The present invention also relates to processes for preparing the compound of formula I. Throughout the following description of such processes it is understood that, where appropriate, suitable protecting groups will be added to, and subsequently removed from, the various reactants and intermediates in a manner that will be readily understood by one skilled in the art of organic synthesis. Conventional procedures for using such protecting groups as well as examples of suitable protecting groups are described, for example, in

"Protective Groups in Organic Synthesis" T. W. Greene, Wiley-Interscience, New York, 1991.

Methods of Preparation of Intermediates

1. In the case where Y is NR₂CO and X is N


(i) Benzoylation of the compound of the formula II, described in: Thorberg S.-O.; Hall H.; Åkesson C.; Svensson K.; Nilsson J. L. G. *Acta Pharm. Suec.* **1987**, *24(4)*, 169-182 as a racemate or in the patent application WO 93/07135 as an enantiomer,



(II)

to obtain a compound of formula III may be carried out by reaction with a suitable benzoylation agent, e.g. a benzyl halide such as benzyl bromide or benzyl chloride or an activated alcohol, e.g. benzyl mesylate or benzyl tosylate. The reaction may be carried out using a salt or the base of compound II in a suitable solvent, e.g. *N,N*-dimethylformamide, acetone or acetonitrile, with a suitable base, e.g. NaOH, NaHCO₃, K₂CO₃ or a trialkylamine such as triethylamine, at a temperature within the range of +20 °C to +150 °C. The presence of a suitable catalyst, e.g. potassium iodide or sodium iodide, may increase the speed of the reaction.

(ii) Demethylation of the compound of formula III




(III)

15

to obtain a compound of formula IV may be carried out by treating the compound with an acidic reagent such as aqueous HBr, HI, HBr//CH₃COOH, BBr₃, AlCl₃, pyridine-HCl or with a basic nucleophilic reagent such as CH₃C₆H₄S⁻ or C₂H₅S⁻ in a suitable solvent. Suitable solvents may be methylene chloride or chloroform and the reaction may occur between -78 °C and +60 °C.


(iii) Conversion of the compound of formula IV to a compound of formula V



10 (IV)

(V)

may be carried out by the reaction with a compound of formula VI




(VI)

where Lg denotes for a leaving group, e.g. a halogen such as chlorine, bromine or iodine or an alkane- or arnesulfonyloxy group such as a p-toluenesulfonyloxy group and R_n and R_b are hydrogen or a lower alkyl group, e.g. methyl. The process may be carried out with a salt of the compound of formula IV obtained by reaction with a base such as K₂CO₃, Na₂CO₃, KOH, NaOH, BuLi or NaH. The reaction may be conducted in a suitable solvent e.g. an aprotic solvent such as dioxane, *N,N*-dimethylformamide, tetrahydrofuran, toluene, benzene or petroleum ether and the reaction may occur between +20 °C and +150 °C.

16

(iv) Rearrangement of a compound of formula V to a compound of formula VII



(V)


(VII)

may be carried out in a suitable solvent, e.g. aprotic solvent such as *N,N*-dimethylformamide, dioxane, 1,1,3,3-tetramethylurea, tetrahydrofuran or hexamethylphosphoric triamide, with a suitable base, e.g. K₂CO₃, KOH, potassium *tert*-butoxide or NaH, at a temperature within the range of +20 °C to +150 °C. The presence of a cosolvent such as 1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone or hexamethylphosphoric triamide in appropriate concentration in the solvent may increase the speed of the reaction.

(v) Hydrolysis of a compound of formula VII to a compound VIII may be carried out under acidic conditions using acids such as H₂SO₄, HCl or HBr in a suitable solvent, e.g. H₂O, ethanol, methanol or mixtures thereof, and the reaction may occur between +20 °C and +100 °C or under basic conditions using bases such as NaOH or KOH in a suitable solvent, e.g. H₂O, ethanol, methanol or mixtures thereof, and the reaction may occur between +20 °C and +100 °C.


17

(vi) Conversion of compound of formula VIII to a compound of formula IX



may be carried out by

a) reaction with a compound of formula X




where R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl. The process may be carried out in a suitable solvent, e.g. an aprotic/anhydrous solvent such as tetrahydrofuran or *N,N*-

dimethylformamide, in the presence of coupling reagent such as *N,N'*-carbonyldiimidazole and the reaction may occur between +20 °C and +130 °C. The reaction is followed by the reduction of the imide with a suitable reducing agent, e.g. $LiAlH_4$, in a suitable solvent, e.g. diethyl ether or tetrahydrofuran at a temperature between +20 °C and reflux, or

20


18

b) by reaction with a compound of formula XI



where LG denotes a leaving group, e.g. a halogen such as chlorine, bromine or iodine or an alkane- or arnesulfonyloxy group such as *p*-toluenesulfonyloxy group and R_1 is hydrogen, C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl. The process may be carried out in a suitable solvent such as ethanol, butanol, *N,N*-dimethylformamide, acetonitrile or a mixture of water and acetonitrile with a suitable base, e.g. K_2CO_3 , $NaHCO_3$ or KOH , and the reaction may occur between +20 °C and +150 °C.

(vii) Conversion of the compound of formula IX to a compound of formula XII



where R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl may be carried out by


a) hydrogenation using a catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent, e.g. acetic acid or ethanol, and at a reaction temperature between +20 °C and +120 °C, or

20

19

b) debenzylolation in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C and at a reaction temperature between +20 °C and reflux.

(viii) Conversion of a compound of formula IX, where R₁ is hydrogen, to a compound of formula XIII,



(IX)

(XIII)

where R_C denotes a suitable protecting group, may be carried out by

a) hydrogenation using a catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent, e.g. acetic acid or ethanol, at a reaction temperature between +20 °C and +120 °C, or


b) debenzylation in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C at a reaction temperature between +20 °C and reflux.

Said reaction is followed by the protection of the piperazine ring in a suitable solvent, e.g. methylene chloride or chloroform, with an appropriate protecting reagent e.g. di-*tert*-butyl dicarbonate with a suitable base, e.g. triethylamine or K₂CO₃, and at a temperature between -20 °C and +60 °C, resulting in compound of formula XIII.

20

2. In the case where Y is NR₂CO and X is CH


(i) Halogenation of the compound of formula XIV, either as a racemate (described in: Thorberg S-O.; Hall H.; Åkesson C.; Svensson K.; Nilsson J. L. G. *Acta Pharm. Suec.* 1987, 24(4), 169-182), or as an enantiomer



(XIV)

(XV)

to obtain a compound of formula XV may be performed by aromatic electrophilic substitution using a suitable halogenation agent such as Br₂, Cl₂, I₂, ICl, or SO₂Cl₂. The reaction may be carried out using the salt or the base of the compound XIV in an appropriate solvent, e.g. acetic acid, HCl/ethanol or water, with or without a suitable base, e.g. an alkali metal acetate such as sodium acetate and at a reaction temperature between -20 °C and room temperature.




(XV)

(ii) Benzylolation of the compound of the formula XV, either as a racemate or as an enantiomer, to obtain a compound of the formula XVI may be carried out by reaction with a suitable benzylolation agent, e.g. benzyl halide such as benzyl bromide or benzyl chloride. The reaction may be carried out using the salt or the base of compound XV in a suitable

21

solvent, e.g. *N,N*-dimethylformamide, acetone or acetonitrile, with a suitable base such as triethylamine, NaOH, NaHCO₃ or K₂CO₃ at a temperature within the range of +20 °C to +150 °C. The presence of a suitable catalyst, e.g. an alkali metal halide such as potassium iodide or sodium iodide, may increase the speed of the reaction.

5




(XVII)

- (iii) The conversion of the compound of the formula XVI to the compound of the formula XVII, where R₁ is C₁-C₆ alkyl or C₃-C₆ cycloalkyl, may be performed by a metal-halogen exchange, in an appropriate anhydrous solvent such as tetrahydrofuran or diethyl ether using a suitable alkyl lithium or metal, e.g. butyllithium, lithium or magnesium turnings, followed by treatment with an appropriate piperidone such as *N*-methyl-4-piperidone and a subsequent suitable workup. The reaction may be performed at a reaction temperature within the range of -78 °C to room temperature.

15


22



(XVIII)

- (iv) The compound of the formula XVII may be reduced to the compound of the formula XVIII by treatment with a suitable reducing agent such as sodium borohydride and a protonating agent such as CF₃COOH, CF₃SO₃H or HCOOH in an appropriate solvent such as tetrahydrofuran or diethyl ether. The reaction may be performed at reaction temperature between 0 °C and reflux.

10




(XIX)

- (v) Demethylation of the compound of the formula XVIII to obtain a compound of formula XIX may be performed by treating the compound with an acidic reagent such as aqueous HBr, HI, HBr/acetic acid, BBH₃, AlCl₃, pyridine-HCl or with a basic nucleophilic reagent

15


such as $C_2H_5S^+$ or $CH_3C_6H_4S^+$ in a suitable solvent. Suitable solvents may be methylene chloride or chloroform and the reaction may occur between $-78^\circ C$ and $+60^\circ C$.

5



(vi) Conversion of the compound of formula XIX to a compound of formula XX may be carried out with a compound such as trifluoromethanesulfonic anhydride in a suitable solvent such as methylene chloride or carbon tetrachloride in the presence of a base such as 2,4,6-collidine, triethylamine or pyridine at a reaction temperature within the range of $-78^\circ C$ to room temperature.

10



15

(vii) Conversion of the compound of formula XX to a compound of formula XXI may be performed by

a) hydrogenation using a catalyst such as palladium, platinum, rhodium or nickel in a suitable solvent such as acetic acid or ethanol at a reaction temperature between $+20^\circ C$ and $+120^\circ C$, or


b) reaction in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C at a reaction temperature between $+20^\circ C$ and reflux.

5

3. In the case where Y is $CONR_2$ and X is N

(i) Nitration of a compound of formula XXII either as a racemate (described in: Thorberg S.-O.; Hall H.; Åkesson C.; Svensson K.; Nilsson J. L. G. *Acta Pharm. Suppl.* 1987, 24(4), 169-182), or as an enantiomer to obtain a compound of formula XXIII.

10




15

where R_d is C_1-C_6 alkyl, may be carried out by aromatic electrophilic substitution using a suitable nitration reagent such as nitric acid or nitric acid and sulfuric acid in a suitable solvent, e.g. acetic acid, acetic anhydride or water, at a reaction temperature between $-20^\circ C$ and room temperature.

20

(ii) Demethylation of the compound of the formula XXIII to obtain a compound of formula XXIV


25



may be carried out by treating the compound with an acidic reagent such as aqueous HBr, HI, HBr/CH₃COOH, BBr₃, AlCl₃, pyridine-HCl or with a basic nucleophilic reagent such as CH₃C₆H₄S⁻ or C₂H₅S⁻. Suitable solvents may be methylene chloride or chloroform and the reaction may occur between -78 °C and +60 °C.

During the demethylation of XXIII, hydrolysis of the ester may occur and the acid function could then be converted back to the ester by methods known by a person skilled in the art (See T. W. Greene, Wiley-Interscience, New York, 1991).


(iii) Conversion of the compound of formula XXIV to a compound of formula XXV



may be carried out by the reaction with an activated trifluoromethanesulfonic reagent e.g. trifluoromethanesulfonic anhydride in a suitable solvent such as methylene chloride, chloroform or carbon tetrachloride in the presence of a suitable base such as triethylamine, pyridine or 2,4,6-collidine at a reaction temperature between -78 °C and room temperature.

26


(iv) Conversion of the compound of formula XXV to a compound of formula XXVI may be carried out by



a) hydrogenation using a catalyst containing palladium, platinum or nickel in a suitable solvent such as ethanol, methanol or acetic acid and at a reaction temperature between +20 °C and +120 °C or


b) reaction in a suitable solvent such as methanol in the presence of an ammonium formate such as triethyl ammonium formate and Pd/C and at a reaction temperature between +20 °C and reflux.

(v) Conversion of the compound of formula XXVI to a compound of formula XXVII




may be carried out by reaction of compound XI

27




where Lg denotes a leaving group, e.g. a halogen such as chlorine, bromine or iodine, or an alkane- or arenesulfonyloxy group such as p-toluenesulfonyloxy group and R₁ is hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl. The process may be carried out in a suitable solvent such as ethanol, butanol, *N,N*-dimethylformamide, acetonitrile or a mixture of water and acetonitrile with a suitable base, e.g. K₂CO₃, NaHCO₃ or KOH, and the reaction may occur between +20 °C and +150 °C. During the cyclization reaction of XXVI, hydrolysis of the ester may occur.



(vi) Hydrolysis of a compound of formula XXVII may be carried out under acidic conditions using acids such as H₂SO₄, HCl, HBr, in a suitable solvent such as H₂O, ethanol, methanol, acetic acid or mixtures thereof at a temperature between + 20 °C and reflux or under basic conditions using bases such as NaOH or KOH in a suitable solvent such as H₂O, ethanol, methanol or mixtures thereof at a temperature between + 20 °C and reflux, resulting in a compound of formula XXVIII, where R₁ is hydrogen, C₁-C₆ alkyl or C₃-C₆ cycloalkyl.


28

(vii) When R₁ is hydrogen, protection of a compound of formula XXVIII as a compound of formula XXIX where R_c is a protecting group



may be carried out by the reaction with a suitable protecting reagent such as di-*tert*-butyl dicarbonate in a suitable solvent, e.g. methylene chloride or chloroform, with a suitable base such as triethylamine or K₂CO₃ and at a temperature between - 20 °C and + 60 °C.

(i) Conversion of a compound of formula XXX to a compound of formula XXXI




may be carried out by

a) hydrolysis of the nitrile in compound of formula XXX in a suitable solvent such as aqueous methanol or aqueous ethanol in the presence of a suitable base such as NaOH or KOH at a reaction temperature between room temperature and reflux, followed by

29


b) hydrolysis of the above formed amide and the ketal under acidic conditions in a suitable solvent such as aqueous methanol, aqueous ethanol or water in the presence of a suitable acid such as HCl or HBr at a reaction temperature between room temperature and reflux.

5 (ii) Conversion of a compound of formula XXXI to a compound of formula XXXII




may be carried out by reaction with a suitable azide such as sodium azide in a suitable acid or mixtures of acids such as H_2SO_4 and acetic acid at a reaction temperature between 0 °C and +50 °C.

(iii) Conversion of a compound of formula XXXIII to a compound of formula XXXIV




may be carried out by reaction with 1-benzylpiperazin in a suitable solvent such as *N,N*-dimethylformamid, dimethylsulfoxid or acetonitril in the presence of a suitable base such as KOH or K_2CO_3 at a reaction temperature between +50 °C and +150 °C.



23 (iv) Hydrolysis of a compound of formula XXXIV to a compound XXXV may be carried out under acidic conditions using acids such as H_2SO_4 , HCl or HBr in a suitable solvent.


30

e.g. H_2O , ethanol, methanol or mixtures thereof, and the reaction may occur between +20 °C and +100 °C or under basic conditions using bases such as NaOH or KOH in a suitable solvent, e.g. H_2O , ethanol, methanol or mixtures thereof, and the reaction may occur between +20 °C and +100 °C.



5 (v) Halogenation of a compound of formula XXXVI to a compound of formula XXXVII where Hal denotes bromine, chlorine or iodine may be performed by a reagent such as [C] or Br_2 , Cl_2 or SO_2Cl_2 with a suitable base such as sodium acetate in a suitable solvent such as acetic acid at a reaction temperature between +20 °C and +50 °C.

(vi) Conversion of a compound of formula XXXVII to a compound of formula XXXVIII



may be carried out by a metal-halogen exchange, in an appropriate anhydrous solvent such as tetrahydrofuran or diethyl ether using a suitable alkyl-lithium or metal, e.g. butyllithium, lithium or magnesium turnings, followed by treatment with carbon dioxide at a reaction temperature between -78 °C and room temperature.


Methods of Preparation of End Products

Another object of the invention is a process A(i), A(ii), A(iii), B(i), B(ii) or C for the preparation of the compound of general formula I by

31

A(ii)

acylation, in the case when R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, Y is NR_2CO , R_2 is hydrogen and X and R_3 are as defined in general formula I above, of a compound of formula A,




- 10 with an activated carboxylic acid $R_3-CO-Lg_1$, where Lg_1 is a leaving group or by using a carboxylic acid R_3-COOH with an activating reagent.

Thus, the acylation according to the process A(i) may be carried out with an appropriate activated carboxylic acid, R_3CO-Lg_1 , where R_3 is as defined above and Lg_1 is a leaving group, such as halogen, e.g. chlorine, in a suitable solvent such as methylene chloride or chloroform with a suitable base, e.g. a trialkylamine such as triethylamine, at a temperature between $-20^\circ C$ and reflux temperature or by using an carboxylic acid, R_3COOH wherein R_3 is as defined above with an activating reagent, e.g. N,N' -carbonyldiimidazole, N,N' -dicyclohexylcarbodiimide or diphenylphosphinic chloride, with a suitable base such as N -methylmorpholine in a suitable solvent such as N,N -dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature between $+20^\circ C$ and $+150^\circ C$.

A(iii)

acylation, in the case when R_1 is hydrogen, Y is NR_2CO , R_2 is hydrogen, R_3 is a protecting group and X and R_3 are as defined in general formula I above, of a compound of formula B

32




- 10 with an activated carboxylic acid $R_3-CO-Lg_1$, where Lg_1 is a leaving group or by using a carboxylic acid R_3-COOH with an activating reagent, followed by the removal of the protecting group R_c .

Thus, the acylation according to the process A(ii) may be carried out with an appropriate activated carboxylic acid, R_3CO-Lg_1 , where R_3 is as defined above and Lg_1 is a leaving group, such as halogen, e.g. chlorine, in a suitable solvent such as methylene chloride or chloroform with a suitable base, e.g. a trialkylamine such as triethylamine, or by using a carboxylic acid, R_3COOH where R_3 is defined as above, with an activating reagent, e.g. N,N' -carbonyldiimidazole, N,N' -dicyclohexylcarbodiimide or diphenylphosphinic chloride, with a suitable base such as N -methylmorpholine in a suitable solvent such as N,N -dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature between $+20^\circ C$ and $+150^\circ C$, followed by removal of the protecting group R_c by hydrolysis in a suitable solvent such as methylene chloride or chloroform with a suitable acid such as trifluoroacetic acid at a temperature between $+20^\circ C$ and $+60^\circ C$.

A(iv)

debenzylation, in the case when R_1 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl, X and R_3 is as defined in general formula I above and R_3 below is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, $(CH_2)_mOH$ wherein m is 2-6 or COR_8 , of a compound of formula Ia, followed by

a) hydrogenation, b) alkylation and removal of a protecting group or d) acylation;



Thus, in the case when R_9 is H the hydrogenation a) above of a compound of formula Ia may be carried out by using a catalyst such as palladium, platinum, rhodium or nickel in a suitable solvent such as acetic acid or ethanol at a reaction temperature between +20 °C and +120 °C, or reaction in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C at a reaction temperature between +20 °C and reflux.

In the case when R_9 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl the debenzoylation is followed by the alkylation b) above using a suitable alkylation reagent such as R_1 -Lg where Lg is a suitable leaving group, e.g. a halogen such as chlorine, bromine or iodine, or an alkane- or arnesulfonyloxy group such as a p-toluenesulfonyloxy group and R_1 is C_1 - C_6 alkyl. The reaction may be carried out in a suitable solvent such as *N,N*-dimethylformamide, acetone, acetonitrile or tetrahydrofuran with a suitable base such as K_2CO_3 , $NaHCO_3$, NaOH or a trialkylamine such as triethylamine. The reaction may be conducted at a temperature between +20 °C and +120 °C or, reductive alkylation with a compound R_1 -CHO, where R_1 is hydrogen or C_1 - C_3 alkyl, or with a C_3 - C_6 cyclic ketone, in the presence of a reductive agent such as sodium cyanoborohydride, sodium borohydride or catalytically with H_2 and a suitable catalyst containing palladium, platinum, rhodium or nickel in a suitable solvent, e.g. tetrahydrofuran, dioxane, methanol or ethanol. A proton donor such as p-toluenesulfonic acid can be used to catalyze the formation of the imine/enamine and

adjustment of pH to slightly acidic by an appropriate acid such as acetic acid may speed up the reaction.


In the case when R_9 is $(CH_2)_mOH$ and m is 2-6, the debenzoylation is followed by the alkylation c) above by using a suitable alkylation reagent such as $BnO(CH_2)_mLg$ where Lg is a suitable leaving group, e.g. a halogen such as chlorine, bromine or iodine, or an alkane- or arnesulfonyloxy group such as a p-toluenesulfonyloxy group and R_1 is C_1 - C_6 alkyl. The reaction may be carried out in a suitable solvent such as *N,N*-dimethylformamide, acetone, acetonitrile or tetrahydrofuran with a suitable base such as K_2CO_3 , $NaHCO_3$, NaOH or a trialkylamine such as triethylamine and may be conducted at a temperature between +20 °C and +120 °C. The reaction is followed by removal of a protecting group, such as a benzyl group, by hydrogenation using a catalyst such as palladium, platinum, rhodium or nickel in a suitable solvent such as acetic acid or ethanol at a reaction temperature between +20 °C and +120 °C, or reaction in a suitable solvent such as methanol in the presence of ammonium formate and Pd/C at a reaction temperature between +20 °C and reflux.

In the case when R_9 is COR, the debenzoylation is followed by the acylation d) above by using an appropriate activated carboxylic acid, R_4CO_2Lg , where R_4 is as defined above and Lg, is a leaving group, such as halogen, e.g. chlorine, in a suitable solvent such as methylene chloride, chloroform or *N,N*-dimethylformamide with a suitable base, e.g. a trialkylamine such as triethylamine or by using a carboxylic acid, R_4COOH where R_4 is defined as above, with an activating reagent, e.g. *N,N'*-carbonyldiimidazole, *N,N'*-dicyclohexylcarbodiimide or diphenylphosphinic chloride, with a suitable base such as *N*-methylmorpholine in a suitable solvent such as *N,N*-dimethylformamide or tetrahydrofuran and the reaction may be conducted at a temperature between +20 °C and +150 °C.

35

B(i)

reacting, in the case when R_1 is C_1-C_6 alkyl or C_3-C_6 cycloalkyl, Y is $CONR_2$, X , R_2 and R_3 are as defined in general formula I above, an activated carboxylic acid of a compound of formula C,




10 with an aniline or amine HNR_2R_3 .

Thus, conversion according to the process B(i) of a compound of formula C may be carried out by activation of the acid function of a compound as an acid halide such as an acid chloride or by using an activating reagent such as *N,N'*-carbonyldiimidazole or *N,N'*-dicyclohexylcarbodiimide in a suitable solvent, e.g. methylene chloride, chloroform, toluene, *N,N*-dimethylformamide, dioxane or tetrahydrofuran, followed by the addition of an appropriate amine or aniline HNR_2R_3 and the reaction may occur between 0 °C and + 120 °C.

B(ii)

reacting, in the case when R_1 is hydrogen, Y is NR_2CO , R_2 is a protecting group and X , R_2 and R_3 are as defined in general formula I above, an activated carboxylic acid of a compound of formula D

36



5 with an aniline or amine HNR_2R_3 , followed by removal of the protecting group R_2 .


Thus, conversion according to the process B(ii), of a compound of formula D, may be carried out by activation of the acid function of a compound as an acid halide such as an acid chloride or by using an activating reagent such as *N,N'*-carbonyldiimidazole or *N,N'*-dicyclohexylcarbodiimide in a suitable solvent, e.g. methylene chloride, chloroform, toluene, *N,N*-dimethylformamide, dioxane or tetrahydrofuran, followed by the addition of an appropriate amine or aniline HNR_2R_3 and the reaction may occur between 0 °C and + 120 °C, followed by removal of the protecting group R_2 by methods known by a person skilled in the art such as hydrolysis in a suitable solvent such as methylene chloride or chloroform with a suitable acid, e.g. trifluoroacetic acid, at a temperature between + 20 °C and + 60 °C.

C

reaction, in the case when R_1 is C_1-C_6 alkyl or C_3-C_6 cycloalkyl, Y is NR_2CONR_2 , R_2 is hydrogen and X and R_3 are as defined in general formula I above, a compound of formula A,

25

37




5 with a suitable azide in the presence of carboxylic acid, $R_3\text{COOH}$.

Thus, reaction according to the process C may be carried out with an appropriate azide such as diphenylphosphoryl azide in the presence of a carboxylic acid, $R_3\text{COOH}$ where R_3 is as defined above in a suitable solvent such as acetonitrile and the reaction may be conducted at a temperature between $+20^\circ\text{C}$ and reflux temperature.

Intermediates

Another object of the invention is a compound having the formula



wherein

$X = \text{N}$ or CH ;

$Z = \text{NH}_2$ or COOH ;

R_1 is H , $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl.

38

The invention is illustrated but not restricted to the following working examples.

Working Examples

Example 1

(*R*)-3-*N,N'*-Dibenzylamino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran.

(*R*)-3-Amino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran (2.6 g, 14 mmol), K_2CO_3 (7.0 g, 51 mmol), benzylbromide (6.0 g, 35 mmol) and a catalytic amount of potassium iodide were mixed in acetonitrile (100 mL) under nitrogen. The reaction mixture was refluxed for 72 h. The solvent was removed *in vacuo*, and the residue was partitioned between diethyl ether and a 2 M NH_3 solution. The layers were separated, and the aqueous phase was extracted twice with diethyl ether. The ethereal layers were combined and dried (MgSO_4). The solvent was removed *in vacuo* to give a yellow oily residue which was purified by flash chromatography on silica gel (eluent: methylene chloride) affording 3.2 g (64% yield) of the title compound: EIMS (70eV) m/z (relative intensity) 359 (91, M^+). The HCl salt was precipitated from diethyl ether at 0°C and then recrystallized from ethanol/diethyl ether. The crystals were hygroscopic and started to melt at 100°C and melted finally between 118 and 120°C ; $[\alpha]_D^{25} -20^\circ$ (c 0.3, methanol).

Example 2

(*S*)-3-*N,N'*-Dibenzylamino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran.

The title compound was synthesized according to the procedure described for its corresponding (*R*)-enantiomer: $[\alpha]_D^{25}$ (measured on the free base) $+1.6^\circ$ (c 1.0, chloroform).

Example 3

(*R*)-3-*N,N'*-Dibenzylamino-5-hydroxy-3,4-dihydro-2*H*-1-benzopyran.

(*R*)-3-*N,N'*-Dibenzylamino-5-methoxy-3,4-dihydro-2*H*-1-benzopyran hydrochloride (1.6 g, 4.0 mmol) was dissolved in methylene chloride (40 mL) under nitrogen, and the solution

was cooled to -70 °C. A solution of boron tribromide (1.8 g, 7.3 mmol) in methylene chloride (25 mL) was added dropwise over 5 min. The temperature was then allowed to slowly reach 0 °C, and the reaction was stirred overnight. The reaction mixture was carefully poured into a saturated NaHCO₃ solution with stirring. The layers were separated and the aqueous phase was extracted three times with methylene chloride. The organic layers were combined and dried (MgSO₄). The solvent was removed *in vacuo* to give a brownish oily residue which was purified by flash chromatography on silica gel (eluent: methylene chloride) affording 0.14 g (98% yield) of the title compound: $[\alpha]_D^{25}$ -94° (c 0.1, methanol); EIMS (70eV) m/z (relative intensity) 345 (100, M⁺).

Example 4

(*S*)-3-*N,N*-Dibenzylamino-5-hydroxy-3,4-dihydro-2*H*-1-benzopyran.

The title compound was synthesized according to the procedure described for its corresponding (*R*)-enantiomer: $[\alpha]_D^{25}$ +109° (c 1.0, chloroform).

Example 5

(*R*)-2-(3-*N,N*-Dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yloxy)-2-

methylpropanamide.

(*R*)-3-*N,N*-Dibenzylamino-5-hydroxy-3,4-dihydro-2*H*-1-benzopyran (35.4 g, 100 mmol) was dissolved in anhydrous 1,4-dioxane (350 mL) under nitrogen. A dispersion of sodium hydride (60-65% in oil, 5.33 g, 130 mmol) was added in portions. The mixture was stirred for 2 h at room temperature. 2-Bromo-2-methylpropanamide (17.9 g, 110 mmol; described in Coutts, I. G. C.; Southcott, M. R. *J. Chem. Soc. Perkin Trans. 1* 1990, 767-771) was added to the dark greenish solution and was heated at reflux with stirring for 3 h. After cooling, a small amount of water was added, the solution was decanted, and the solvent was removed *in vacuo*. The residue was partitioned between ethyl acetate (350 mL) and a saturated NaHCO₃ solution (50 mL). The organic layer was dried (MgSO₄), and the solvent was removed *in vacuo* to give a brownish residue which was chromatographed on a short column of silica gel (eluent: hexane/ethyl acetate; 55:45) affording 27.6 g (64% yield)

of the title compound as a white solid: mp 132-134 °C; $[\alpha]_D^{25}$ -92° (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 430 (6, M⁺).

Example 6

(*S*)-2-(3-*N,N*-Dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yloxy)-2-

methylpropanamide.

The title compound was synthesized according to the procedure described for its corresponding (*R*)-enantiomer: $[\alpha]_D^{25}$ +99° (c 1.0, chloroform).

Example 7

(*R*)-5-Amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran.

1,3-Dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone (31 mL) was added to a stirred solution of (*R*)-2-(3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yloxy)-2-methylpropanamide (31.0 g, 72.0 mmol) in anhydrous *N,N*-dimethylformamide (310 mL) under nitrogen. Sodium hydride (60-65% in oil, 5.76 g, 144 mmol) was added in portions. The reaction mixture was heated at 100 °C and was stirred for 16 h. The mixture was then allowed to cool, and the solution was partitioned between ethyl acetate (500 mL) and a 2 M NH₃ solution (300 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (150 mL). The combined organic layers were dried (MgSO₄) and concentrated *in vacuo* to give a brownish oil. The obtained material was dissolved in ethanol (400 mL). A 6 M HCl solution (500 mL) was added, and the reaction mixture was heated to reflux at 85 °C. After stirring overnight, the mixture was allowed to cool to 35 °C, the ethanol solvent was concentrated *in vacuo*, and toluene was added to the residual aqueous solution. The mixture was cooled on ice, and a solution of conc. NH₃ was slowly added with stirring. An almost insoluble material formed. The alkaline two-phase system was transferred to a separatory funnel, and the insoluble material was treated with a 2 M NH₃ solution and ethyl acetate. Eventually, all material was dissolved and it was combined with the already obtained two-phase mixture. The layers were separated, and the aqueous layer was extracted with another portion of ethyl acetate. The combined organic layers

were dried ($MgSO_4$), and the solvent was removed *in vacuo* to give a brownish oil which was purified on a short column of silica gel (eluent: hexane/ethyl acetate: 80:20) affording 19.0 g (72% yield) of the desired compound as a light yellow oil. The product slowly crystallized upon standing in the refrigerator: mp 99-101 °C; $[\alpha]_D^{25}$ -131° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 344 (38, M^+).

Example 8

(*S*)-5-Amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran.

The title compound was synthesized according to the procedure described for its corresponding (*R*)-enantiomer: $[\alpha]_D^{25}$ +123° (c 1.0, chloroform). An analytical sample was recrystallized from diethyl ether/petroleum ether: mp 101-103 °C.

Example 9

(*R*)-3-*N,N*-Dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran.

To a solution of (*R*)-5-amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran (2.86 g, 8.30 mmol) in a mixture of 15% water in acetonitrile (120 mL) were added sodium iodide (69 mg, 0.42 mmol) and *N*-methyl-bis(2-chloroethyl)amine hydrochloride (3.20 g, 16.6 mmol) with stirring. The clear solution was heated at reflux. After 7 h of stirring, $NaHCO_3$ (700 mg, 8.30 mmol) was added, and the reaction mixture was stirred for an additional 11 h. Another portion of $NaHCO_3$ (700 mg, 8.30 mmol) was added followed by continued reflux. After 6 h, a final portion of $NaHCO_3$ (350 mg, 4.15 mmol) was added, and the reaction mixture was stirred for 6 h more (30 h in all). The mixture was cooled on an ice-bath, and a 2 M NaOH solution (20 mL) was added with stirring. The two-phase system was stirred for 10 min after which the solvents were removed under reduced pressure until a precipitation occurred. The aqueous residue was extracted with diethyl ether (150 mL), the layers were separated, and the aqueous layer was extracted with diethyl ether (2 x 50 mL). The combined ethereal layers were dried ($MgSO_4$), and the solvent was removed *in vacuo*. The crude product was purified by column chromatography on silica (eluent: chloroform/ethanol: 95:5:4.5 + 0.5% conc. NH_3) affording 2.39 g (67% yield) of the title

compound as a colorless oil: $[\alpha]_D^{25}$ -45° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 427 (0.3, M^+).

Example 10

(*S*)-1-(3-*N,N*-Dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yl)-4-methylpiperazine-2,6-dione.

To a dispersion of *N*-methylimidodiacetic acid (6.90 g, 46.9 mmol) in anhydrous tetrahydrofuran (575 mL) was added 1,1'-carbonyldiimidazole (15.2 g, 93.9 mmol), and the mixture was heated at reflux for 2 h under nitrogen. A solution of (*S*)-5-amino-3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran (15.0 g, 42.7 mmol) in tetrahydrofuran (120 mL) was added with stirring over 0.5 h. The reaction mixture was heated at reflux for 28 h, then allowed to cool, and the solvent was removed *in vacuo*. The residue was purified on a short column of silica gel (eluent: methylene chloride and ethyl acetate) affording 14.1 g (71% yield) of the title compound as a light yellow solid: mp 59-60 °C; $[\alpha]_D^{25}$ +89° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 455 (8, M^+).

Example 11

(*S*)-3-*N,N*-Dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2*H*-1-benzopyran.

To a stirred solution of (*S*)-1-(3-*N,N*-dibenzylamino-3,4-dihydro-2*H*-1-benzopyran-5-yl)-4-methylpiperazine-2,6-dione (25.4 g, 55.8 mmol) in anhydrous diethyl ether (800 mL) was added lithium aluminum hydride (9.30 g, 246 mmol) in portions. The reaction mixture was heated to reflux for 6.5 h under nitrogen and was stirred overnight at room temperature. The mixture was cooled (ice-bath), and water (10 mL) was added followed by a 15% aqueous solution of NaOH (10 mL) and another portion of water (30 mL). The precipitate was filtered off and washed with several portions of warm tetrahydrofuran. The organic layers were combined, and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol: 95:5 + 0.5% conc. NH_3) affording 13.6 g (57% yield) of the title compound as a light yellow oil: $[\alpha]_D^{25}$ +63° (c 1.0, methanol); EIMS (70eV) *m/z* (relative intensity) 427 (3, M^+).

Example 12

(R)-3-Amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran.

To a solution of (R)-3-*N,N*-dibenzylamino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (2.34 g, 5.47 mmol) in anhydrous methanol (100 mL) were added palladium (10%) on activated carbon (0.86 g) and ammonium formate (2.76 g, 43.8 mmol) under nitrogen. The reaction mixture was heated at 50 °C with stirring overnight. The solution was filtered through Celite®, and the solvent was removed *in vacuo*. The residue was partitioned between a 2 M NH₃ solution (20 mL) and ethyl acetate (100 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (3 x 50 mL). The combined organic phases were dried (Na₂SO₄), and the solvent was removed *in vacuo* to give 1.21 g (90% yield) of the title compound as a pale yellow oil: [α]_D²⁵ +15° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 247 (6, M⁺).

Example 13

(S)-3-Amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran.

The title compound was synthesized according to the procedure described for its corresponding (R)-enantiomer: [α]_D²⁵ -15° (c 1.0, chloroform).

Example 14

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-

morpholinobenamide.

A solution of 4-morpholinobenzoic acid (380 mg, 1.83 mmol; described in: Deguis, J.; Rastickiene, L.; Degutene, A. *Zh. Orq. Khim.* 1978, 14(10), 2060-2064) and 1,1'-

carbonyldiimidazole (310 mg, 1.92 mmol) in anhydrous *N,N*-dimethylformamide (12 mL) was stirred at 75 °C for 30 min. The mixture was allowed to cool after which a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (430 mg, 1.74 mmol) in *N,N*-dimethylformamide (8 mL) was added. The reaction mixture was stirred at room temperature for 3 days. Another portion of 1,1'-carbonyldiimidazole (57 mg, 0.35 mmol) was added, and the mixture was stirred for an additional 3.5 h. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica

(eluent: chloroform/ethanol; 93:7 + 0.5% NH₃) affording 513 mg (68% yield) of the title compound as a white solid: mp 210-212 °C; [α]_D²⁵ -145° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 436 (65, M⁺).

Example 15

(R)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-

morpholinobenamide.

The title compound was synthesized according to the procedure described for its corresponding (S)-enantiomer: [α]_D²⁵ +145° (c 1.0, chloroform).

Example 16

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-

piperidinobenamide.

A suspension of 4-piperidinobenzoic acid (276 mg, 1.35 mmol; described in: Wennga, W. D.; Janssen, M. J. *Recl. Trav. Chim. Pays-Bas* 1968, 87(12), 1372-1380) and 1,1'-carbonyldiimidazole (229 mg, 1.41 mmol) in anhydrous *N,N*-dimethylformamide (11 mL) was placed in an oil bath at 75 °C. After 45 min of stirring, the mixture was allowed to cool. A solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (317 mg, 1.28 mmol) in *N,N*-dimethylformamide (5 mL) was added, and the mixture was stirred at room temperature for 40 h. Another portion of 1,1'-carbonyldiimidazole (83 mg, 0.51 mmol) was added, and the reaction was stirred for 3 days. At this time, the reaction was not complete and a final amount of 1,1'-carbonyldiimidazole (42 mg, 0.25 mmol) was added. The reaction mixture was heated for 3 h at 50 °C after which the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% NH₃) affording 202 mg (36% yield) of the title compound as a white solid: mp 178-180 °C; [α]_D²⁵ -159° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 434 (35, M⁺).

Example 17

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-butoxybenzamide.

A solution of 4-butoxybenzoic acid (650 mg, 3.35 mmol) in thionyl chloride (13 mL) was heated at 50 °C for 15 min after which the mixture was allowed to reach room temperature. The excess of thionyl chloride was removed under reduced pressure, and the residue was evaporated with two portions of toluene. The acid chloride was obtained as a brownish oil. A portion of the acid chloride (150 mg, 0.705 mmol) was dissolved in methylene chloride (5 mL) and added to an ice-cooled solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (159 mg, 0.643 mmol) and triethylamine (134 µL, 0.960 mmol) in anhydrous methylene chloride (20 mL). The ice-bath was removed, and the temperature was allowed to reach room temperature. The reaction mixture was washed with a saturated NaHCO₃ solution (10 mL), dried (MgSO₄), and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% conc. NH₃) affording 200 mg (74% yield) of the title compound as a white solid: mp 192-193.6 °C; [α]_D²² -114° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 423 (52, M⁺).

Example 18

(R)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-butoxybenzamide.

The title compound was synthesized according to the procedure described for its corresponding (S)-enantiomer: [α]_D²¹ +104° (c 1.0, chloroform).

Example 19

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-trifluoromethylbenzamide.

A mixture of 4-trifluoromethylbenzoic acid (195 mg, 1.02 mmol) in thionyl chloride (5 mL) was heated at 50 °C for 20 min and then to reflux for 10 min. The mixture was allowed to cool after which the excess of thionyl chloride was removed *in vacuo*, and the

residue was evaporated with two portions of toluene. The acid chloride was then dissolved in anhydrous methylene chloride (5 mL) and added to an ice-cooled solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (230 mg, 0.930 mmol) and triethylamine (194 µL, 1.39 mmol) in anhydrous methylene chloride (20 mL) with stirring. The reaction mixture was allowed to reach room temperature and was washed with a saturated NaHCO₃ solution. After drying (MgSO₄) and evaporation of the solvent *in vacuo*, a crude product was obtained which was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% conc. NH₃). This procedure gave 214 mg (35 % yield) of the title compound as a white solid: mp 212-214 °C; [α]_D²² -73° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 419 (100, M⁺).

Example 20

(R)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-trifluoromethylbenzamide.

The title compound was synthesized according to the procedure described for its corresponding (S)-enantiomer: [α]_D²¹ +74° (c 1.0, chloroform).

Example 21

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-2,4-dimethoxybenzamide.

A solution of 2,4-dimethoxybenzoic acid (185 mg, 1.01 mmol) in thionyl chloride (5 mL) was heated at 55 °C for 15 min. The excess of thionyl chloride was removed *in vacuo*, and the residue was evaporated with two portions of toluene. The acid chloride was then dissolved in anhydrous methylene chloride (5 mL) and added to an ice-cooled solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (228 mg, 0.920 mmol) in anhydrous methylene chloride (20 mL) with stirring. The precipitated product was dissolved by the addition of triethylamine (193 µL, 1.38 mmol) to give a clear light yellow solution. The ice-bath was removed, and the reaction mixture was stirred at room temperature for 1 h. The mixture was washed with a saturated NaHCO₃ solution, dried (MgSO₄), and the solvent was removed *in vacuo*. The residue was purified by column

chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% conc. NH_3) affording 268 mg (71% yield) of the title compound as an oil: $[\alpha]_D^{25}$ -91° (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 411 (4, M^+). The base (238 mg, 0.578 mmol) was dissolved in anhydrous diethyl ether (10 mL) under nitrogen and was cooled on an ice-bath. A solution of HCl in diethyl ether (3 M, 0.5 mL), diluted with diethyl ether (5 mL), was added dropwise with stirring. The HCl salt was filtered, washed with diethyl ether, and dried *in vacuo* affording 187 mg (69% yield) of the product as a white powder: mp siners >44 °C.

Example 22

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-N,N'-diethylaminobenzamide.

A solution of 4-diethylaminobenzoic acid (189 mg, 0.978 mmol) and 1,1'-carbonyldiimidazole (166 mg, 1.02 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was stirred at 75 °C for 45 min. The mixture was allowed to cool, and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (230 mg, 0.930 mmol) in *N,N*-dimethylformamide (8 mL) was added. The reaction mixture was stirred at room temperature for 7 days. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% NH_3) affording 234 mg (60% yield) of the title compound as a white solid: mp 218-219.6 °C; $[\alpha]_D^{25}$ -178° (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 422 (29, M^+).

Example 23

(R)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-N,N'-diethylaminobenzamide.

The title compound was synthesized according to the procedure described for its corresponding (S)-enantiomer: $[\alpha]_D^{25}$ +172° (c 1.0, chloroform).

Example 24

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-furan-2-carboxamide.

To an ice-cooled stirred solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (230 mg, 0.930 mmol) and triethylamine (194 μL , 1.39 mmol) in anhydrous methylene chloride (10 mL) was added 2-furyl chloride (101 μL , 1.02 mmol) under nitrogen. The ice-bath was removed, and the reaction mixture was allowed to reach room temperature. The mixture was washed with a 2 M NH_3 solution, dried (MgSO_4), and the solvent was removed *in vacuo*. The remains was purified on a chromatotron (accelerated thin layer chromatography, eluent: chloroform/ethanol; 92:8 + 0.5% conc. NH_3) affording 249 mg (79% yield) of the title compound as a white solid: mp siners >50 °C; $[\alpha]_D^{25}$ -83° (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 341 (32, M^+).

Example 25

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-N,N'-dimethylaminobenzamide.

A solution of 4-dimethylaminobenzoic acid (190 mg, 1.15 mmol) and 1,1'-carbonyldiimidazole (205 mg, 1.26 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was stirred at 75 °C for 35 min. The mixture was allowed to cool, and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (271 mg, 1.10 mmol) in *N,N*-dimethylformamide (5 mL) was added. The reaction mixture was stirred at room temperature for 4 days. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% NH_3) affording 292 mg (67% yield) of the title compound as a white solid: mp 248-250 °C; $[\alpha]_D^{25}$ -173° (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 394 (46, M^+).

Example 26

(S)-N-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-pyrrole-2-carboxamide.

A mixture of 1,1'-carbonyldiimidazole (360 mg, 1.85 mmol) and pyrrole-2-carboxylic acid (225 mg, 2.03 mmol) in anhydrous *N,N*-dimethylformamide (8 mL) was stirred at 75 °C for 45 min. The mixture was allowed to cool, and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (457 mg, 1.85 mmol) in *N,N*-dimethylformamide (10 mL) was added. The reaction mixture was stirred for 7 days at room temperature under nitrogen. The solvent was removed *in vacuo*, and the residue was extracted with diethyl ether (50 mL) and water (20 mL). The aqueous layer was extracted with another portion of diethyl ether (50 mL). The combined ethereal layers were dried (MgSO_4), and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 90:10 + 0.5% conc. NH_3) yielding the title compound as an oil. Evaporation with diethyl ether afforded 300 mg (48% yield) of the title compound as a white powder: mp siners >96 °C; $[\alpha]_D^{25}$ -82.8° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 340 (10, M^+).

Example 27

(S)-N-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-5-

methylpyridine-3-carboxamide.

A solution of 5-methylnicotinic acid (141 mg, 1.03 mmol) and 1,1'-carbonyldiimidazole (183 mg, 1.13 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was stirred at 75 °C for 55 min. The mixture was allowed to cool, and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (232 mg, 0.94 mmol) in *N,N*-dimethylformamide (5 mL) was added. The reaction mixture was stirred at room temperature for 28 h. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 87:13 + 0.5% NH_3). The product was contaminated with a large amount of imidazole which could be removed by the following procedure: The mixture was dissolved in diethyl ether (100 mL), washed with water (2 x 20 mL) and treated with brine (10 mL). The ethereal layer was dried

(MgSO_4), and the solvent was removed *in vacuo* to give 119 mg (35% yield) of the title compound as a white solid: mp siners >68 °C; $[\alpha]_D^{25}$ -82° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 366 (21, M^+).

Example 28

(S)-N-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-2,4-bis(trifluoromethyl)benzamide.

A solution of 2,4-bis(trifluoromethyl)benzoic acid (195 mg, 0.755 mmol) in thionyl chloride (4 mL) was heated at 55 °C for 45 min. The excess of thionyl chloride was removed *in vacuo*, and the residue was evaporated with two portions of toluene. The acid chloride was then dissolved in anhydrous methylene chloride (5 mL) and added to a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (170 mg, 0.687 mmol) and triethylamine (144 μL , 1.03 mmol) in anhydrous methylene chloride (20 mL) with stirring. The reaction mixture was left overnight at room temperature and was washed with a 2 M NH_3 solution (10 mL) followed by a portion of brine. The organic layer was dried (MgSO_4), and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5% conc. NH_3) affording 100 mg (30% yield) of the title compound as a white powder: mp 202-203 °C; $[\alpha]_D^{25}$ -51° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 487 (16, M^+).

Example 29

(S)-N-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-2-hydroxy-4-methoxybenzamide.

A solution of 4-methoxy-2-acetoxybenzoic acid (232 mg, 1.10 mmol), described in: Schonhofer, F. *Ber Deutsch Chem Ges* 1951, 84, 13) in thionyl chloride (5 mL) was heated at 55 °C for 30 min. The excess of thionyl chloride was removed *in vacuo*, and the residue was evaporated with two portions of toluene. The acid chloride was then dissolved in anhydrous methylene chloride (5 mL) and added to a stirred solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (248 mg, 1.00 mmol) and triethylamine (210 μL , 1.50 mmol) in anhydrous methylene chloride (20 mL). The reaction

10 mixture was stirred at room temperature for 2.5 h after which the mixture was washed with
a saturated NaHCO_3 solution, dried (MgSO_4), and the solvent was removed *in vacuo*. The
residue was dissolved in absolute ethanol (20 mL), and conc. NH_3 (5 mL) was added. The
mixture was stirred overnight. The solvent was removed *in vacuo*, and the remains was
5 purified by column chromatography on silica (eluent: chloroform/ethanol; 92:8 + 0.5%
conc. NH_3) affording 120 mg (33% yield) of the title compound as a white solid: mp
sinters >80 °C; $[\alpha]_D^{25}$ -92° (c 1.0, chloroform); EIMS (70eV) *m/z* (relative intensity) 397
(27, M^+).

10 Example 30

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-
trifluoromethoxybenzamide.

15 A mixture of 4-trifluoromethoxybenzoic acid (254 mg, 1.23 mmol) in thionyl chloride (5
mL) was heated at 60 °C for 25 min. The excess of thionyl chloride was removed under
reduced pressure, and the remains were evaporated with two portions of toluene. The acid
chloride was then dissolved in anhydrous methylene chloride (5 mL) and added to a stirred
solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (277 mg,
1.12 mmol) and triethylamine (234 μL , 1.68 mmol) in methylene chloride (10 mL). The
20 reaction mixture was stirred for 2 h at room temperature and was washed with a saturated
solution of NaHCO_3 . The organic layer was dried (MgSO_4), and the solvent was removed
in vacuo. The product was purified by column chromatography on silica (eluent:
chloroform/ethanol; 92:8 + 0.5% conc. NH_3) affording 248 mg (31% yield) of the title
compound as a white solid: mp 192-193 °C; $[\alpha]_D^{25}$ -75° (c 1.0, chloroform); EIMS (70eV)
25 *m/z* (relative intensity) 435 (6, M^+).

25 Example 31

4-(4-Piperidon-1-yl)benzoic Acid.

A solution of 2 M NaOH (10 mL), 4-(8-aza-1,4-dioxaspiro[4,5]dec-8-yl)benzonitrile (820
mg, 3.36 mmol; described in: Taylor E. C.; Skonicki J. S. *Synthesis* 1981, 8, 606-608), and
30 ethanol (7.5 mL) was heated at reflux for 3 h. The external heating was interrupted, and the

reaction mixture was stirred overnight at ambient temperature. The ethanolic solvent was
removed *in vacuo*, and the remains were acidified to pH 4 with a 2 M HCl solution
followed by extraction with ethyl acetate (50 mL). The layers were separated, and pH was
adjusted to pH 6 with a 2 M NaOH solution followed by another extraction with ethyl
5 acetate (50 mL). The combined organic layers were concentrated *in vacuo*, and the solid
residue was dissolved in a 6 M HCl solution (10 mL). The reaction mixture was heated at
75 °C for 2.5 h and then at 55 °C overnight. The temperature was raised to 75 °C for 2 h,
and the reaction mixture was then allowed to cool. The pH was adjusted to pH 4, and the
solution was extracted with ethyl acetate (50 mL). The layers were separated, and another
10 extraction was made at pH 5. The combined organic layers were dried (MgSO_4), and the
solvent was removed *in vacuo*. The crude product was recrystallized from ethyl acetate
affording 300 mg (41% yield) of the title compound as yellowish crystals: mp sinters >215
°C; EIMS (70eV) *m/z* (relative intensity) 219 (100, M^+)

15 Example 32

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-piperidon-
1-yl)benzamide.

20 A solution of 1,1'-carbonyldiimidazole (116 mg, 0.716 mmol) and 4-(4-piperidon-1-
yl)benzoic acid (150 mg, 0.683 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was
stirred at 75 °C for 50 min. The mixture was allowed to cool, and a solution of (S)-3-
amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (161 mg, 0.651 mmol) in
N,N-dimethylformamide (4 mL) was added. The reaction mixture was stirred at room
temperature for 8 days. The solvent was removed *in vacuo*, and the residue was purified by
column chromatography on silica (eluent: chloroform/ethanol; 90:10 + 0.5% conc. NH_3)
25 affording 54 mg (19% yield) of the title compound as a white solid: mp 222-225 °C
(decomposes); $[\alpha]_D^{25}$ -136° (c 0.30, chloroform); TSPMS (70eV) *m/z* 449 ($\text{M}+1$).

Example 33

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-morpholinobenzenesulfonamide.

To a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (120 mg, 0.485 mmol) in anhydrous methylene chloride (10 mL) were added triethylamine (81 μ L, 0.582 mmol) and 4-(4-morpholinyl)benzenesulfonyl chloride (140 mg, 0.534 mmol; described in: Galliani, G. Eur. Pat. Appl. EP 335,758, 1989, *Chem. Abstr.* 1990, 112, 98374d [125393-22-8]). The reaction mixture was stirred at room temperature for 4 h, washed with a 2 M NH_3 solution, dried (MgSO_4) and concentrated *in vacuo*. The crude product was purified by column chromatography on silica (eluent: chloroform/ethanol, 90:10 + 0.5% conc. NH_3) affording 141 mg (61% yield) of the title compound as a white solid; mp siniers > 100 °C; $[\alpha]_D^{22} + 10^\circ$ (c 1.0, chloroform); EIMS (70eV) m/z (relative intensity) 472 (56, M^+).

Example 34

4-(Hexahydro-1,4-diazepin-5-on-1-yl)benzoic Acid.

A solution of 4-(piperidin-1-yl)benzoic acid (281 mg, 1.28 mmol), conc. acetic acid (2 mL), and conc. H_2SO_4 (1 mL) was cooled to 5 °C. Sodium azide (92 mg, 1.41 mmol) was added, and the reaction mixture was stirred at 7 °C for 42 h. A solution of 2 M NaOH was added to pH 5, and the resulting precipitate was filtered and washed with several portions of ice-cooled water. Drying *in vacuo* afforded 272 mg (91% yield) of the title compound as a white solid; mp 285-286 °C; EIMS (70eV) m/z (relative intensity) 234 (66, M^+).

Example 35

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(hexahydro-1,4-diazepin-5-on-1-yl)benzamide.

A solution of 1,1'-carbonyldiimidazole (151 mg, 0.934 mmol) and 4-(hexahydro-1,4-diazepin-5-on-1-yl)benzoic acid (219 mg, 0.934 mmol) in anhydrous *N,N*-dimethylformamide (7 mL) was stirred at 75 °C for 55 min. The mixture was allowed to cool, and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-

benzopyran (210 mg, 0.85 mmol) in *N,N*-dimethylformamide (3.5 mL) was added. The reaction mixture was stirred at room temperature for 14 days. The solvent was removed *in vacuo*, and the residue was purified by column chromatography on silica (eluent: chloroform/ethanol, 90:10 + 1% conc. NH_3). The product was crystallized from a mixture of chloroform, ethanol, and ethyl acetate affording 84 mg (21% yield) of the title compound as white crystals; mp 244-247 °C (decomposes); $[\alpha]_D^{21} - 148^\circ$ (c 0.50, chloroform); TSPMS (70eV) m/z 464 ($\text{M}+1$).

Example 36

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-N'-(4-morpholino)phenyl urea.

To a stirred solution of 4-morpholinobenzoic acid (126 mg, 0.606 mmol) and (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (150 mg, 0.606 mmol) in acetonitrile (5 mL) was diphenylphosphoryl azide (131 μ L, 0.606 mmol) added. The reaction mixture was heated at reflux for 1.5 h and was then allowed to cool to room temperature overnight. The solvent was removed *in vacuo*, and the residue was partitioned between ethyl acetate and a 2 M NH_3 solution. The organic layer was dried (MgSO_4) and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol, 90:10 + 0.5% conc. NH_3) affording 100 mg (36% yield) of the title compound as a white solid; mp siniers > 118 °C; $[\alpha]_D^{21} - 71^\circ$ (c 0.5, chloroform); MSTSP 452 ($\text{M}+1$).

Example 37

4-Bromo-3-methoxymorpholinobenzene.

To a stirred slurry of 4-(3-methoxyphenyl)morpholine (1.54 g, 7.97 mmol; described in: Skowronska-Plasinska M.; Verboon W.; Reinhold D. N. *J. Org. Chem.* 1985, 50/15), 2690-8) and sodium acetate (0.784 g, 9.56 mmol) in 1,4-dioxane (100 mL) was added a 0.25 M solution of bromine in 1,4-dioxane (35.0 mL, 8.77 mmol) over 45 min. Another portion of the bromine solution (15.0 mL, 4.00 mmol) and sodium acetate (0.523 g, 6.38 mmol) were added, and the reaction mixture was heated at 50 °C overnight. The solvent

was removed *in vacuo*, and the residue was partitioned between diethyl ether (100 mL) and a 2 M NH₃ solution. The layers were separated, and the aqueous layer was extracted with diethyl ether (50 mL). The combined organic layers were dried (MgSO₄), and the solvent was removed *in vacuo*. The residue was filtered through a column of silica gel (eluent: chloroform/ethanol, 1:1 + 1.5% conc. NH₃), and the solvent was removed *in vacuo*. The residue was partitioned between methylene chloride and a 2 M NH₃ solution. After drying (MgSO₄) of the organic layer and removal of the solvent *in vacuo*, an orange oil was obtained which was purified by column chromatography on silica (eluent: methylene chloride + 0.5% conc. NH₃) affording 450 mg (21% yield) of the title compound as a white solid: mp 103.5-104.5 °C; EIMS (70 eV) *m/z* (relative intensity) 273/271 (56/56, M⁺).

Example 38

2-Methoxy-4-morpholinobenzole Acid.

To a stirred solution of 4-bromo-3-methoxy-1-morpholinobenzene (104 mg, 0.382 mmol) in anhydrous tetrahydrofuran (3 mL) at -78 °C was slowly added *n*-butyl lithium (1.3 M solution in hexanes, 325 µL, 0.420 mmol) under nitrogen. The cooling medium was exchanged with an ice-bath, and the mixture was stirred for 5 min. After cooling again to -78 °C, carbon dioxide from evaporation of dry ice was bubbled through the solution for 10 min. A precipitate was formed, and the reaction mixture was allowed to reach room temperature. Diethyl ether and water were added. The mixture was extracted, the layers were separated, and the aqueous layer was acidified to pH 4. The dark blue aqueous solution was extracted several times with diethyl ether and ethyl acetate at pH 4 to pH 6. The combined organic layers were dried (MgSO₄), and the solvent was removed *in vacuo* affording 60 mg (66% yield) of the title compound as a white solid: mp 158-160 °C; EIMS *m/z* (relative intensity) 237 (100, M⁺).

Example 39

(S)-N-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-2-methoxy-4-morpholinobenzamide.

A stirred solution of 1,1'-carbonyldiimidazole (222 mg, 1.37 mmol) and 2-methoxy-4-morpholinobenzonic acid (176 mg, 0.740 mmol) in anhydrous *N,N*-dimethylformamide (5 mL) was heated at 75 °C for 2 h, and was then allowed to cool. A solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (183 mg, 0.740 mmol) in anhydrous *N,N*-dimethylformamide (4 mL) was added. The reaction mixture was stirred at room temperature for 5 days. The solvent was removed *in vacuo*, and the residue was partitioned between ethyl acetate (50 mL) and a 2 M NH₃ solution (15 mL). The organic layer was dried (MgSO₄), and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica (eluent: chloroform/ethanol, 93:7 + 0.5% conc. NH₃) affording 113 mg (30% yield) of the title compound as an uncolored foam: [α]_D²⁰ -1.41° (c 0.5, chloroform); EIMS (70eV) *m/z* (relative intensity) 466 (20, M⁺).

Example 40

4-(4-Benzylpiperazin-1-yl)benzonitrile.

To a solution of 4-fluorobenzonitrile (3.0 g, 25 mmol) in *N,N*-dimethylformamide (15 mL) were added 1-benzylpiperazine (4.3 mL, 25 mmol) and potassium carbonate (3.4 g, 25 mmol). The reaction mixture was stirred at 120 °C for 13 h. The solvent was evaporated *in vacuo* and the residue was partitioned between ethyl acetate (100 mL) and water (15 mL). The aqueous phase was extracted with ethyl acetate (30 mL), and the combined organic phases were washed twice with brine (10 mL) and dried (MgSO₄). Evaporation of the solvent gave 7.6 g of crude product. Purification of the residue on a silica gel column using ethyl acetate/methylene chloride (1:9) as the eluent afforded 4.0 g (59% yield) of the title compound as a white solid: mp 104-105 °C; EIMS (70eV) *m/z* (relative intensity) 277 (20, M⁺).

Example 41**4-(4-Benzylpiperazin-1-yl)benzoic acid.**

4-(4-Benzylpiperazin-1-yl)benzonitrile (4.0 g, 15 mmol) was dissolved in glacial acetic acid (40 mL), 6 M hydrochloric acid (50 mL) was added and the reaction mixture was stirred at 100 °C for 17 h. The solvent was evaporated, the residue was suspended in water (10 mL) and the pH was adjusted to 3 by addition of 2 M sodium hydroxide (35 mL). The slurry was stirred at 50 °C for 2 h, cooled and the precipitate was filtered and dried *in vacuo* to give 4.1 g of a crude product. The solid was partitioned between methylene chloride (40 mL) and water (220 mL) with 2 M sodium hydroxide (8 mL). The aqueous phase was washed with methylene chloride (40 mL) and the pH was adjusted to 5 with 2 M hydrochloric acid. The aqueous phase was cooled, the precipitate was filtered and dried *in vacuo* to give 1.6 g (38% yield) of the title compound: mp 226 °C (dec); EIMS (70 eV) *m/z* (relative intensity) 296 (44, M⁺).

Example 42**(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-benzylpiperazin-1-yl)benzamide**

A suspension of 4-(4-benzylpiperazin-1-yl)benzoic acid (1.3 g, 4.2 mmol) and 1,1'-carbonyldiimidazole (740 mg, 4.2 mmol) in *N,N*-dimethylformamide (30 mL) was heated to 75 °C for 1.5 h. The reaction mixture was cooled to 50 °C and a solution of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (1.0 g, 4.0 mmol) was added. The solution was stirred at 50 °C for 20 h and the solvent was evaporated *in vacuo* giving 3.5 g of a crude product. Purification by chromatography on a silica gel column using chloroform/ methanol/ concentrated ammonia 95:5:0.5 as the eluent gave 1.7 g (80% yield) of the title compound as a pale yellow solid : mp siniers > 85 °C; TSPMS *m/z* (relative intensity) 526 (100, M⁺); [α]_D²⁵ -130° (c 1.0, chloroform).

Example 43**(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(piperazin-1-yl)benzamide**

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-benzylpiperazin-1-yl)benzamide (1.7 g, 3.2 mmol) was dissolved in methanol (100 mL). Palladium (10%) on activated carbon (510 mg) and ammonium formate (1.6 g, 26 mmol) were added and the reaction mixture was stirred at 50 °C for 19 h. The catalyst was filtered off and the solvent was evaporated *in vacuo* to give 1.3 g (92% yield) of the title compound as a pale yellow solid: mp > 102 °C siniers; EIMS (70eV) *m/z* (relative intensity) 435 (8, M⁺); [α]_D²⁵ -102° (c 0.15, chloroform).

Example 44**(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-acetyl piperazin-1-yl)benzamide**

(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(piperazin-1-yl)benzamide (460 mg, 1.0 mmol) was dissolved in *N,N*-dimethylformamide (5 mL) and acetyl chloride (82 µL, 1.2 mmol) was added.

The solution was stirred at ambient temperature for 1 h and the solvent was evaporated *in vacuo*. The residue was partitioned between methylene chloride (80 mL) and 2 M NaOH (10 mL). The organic layer was washed with brine (5 mL) and dried (MgSO₄). Evaporation of the solvent *in vacuo* gave 660 mg of a crude product. Purification by column chromatography on silica using chloroform/ethanol (saturated with ammonia) 15:1 as the eluent afforded 330 mg (66% yield) of the title compound as a white solid: mp 88 °C (dec); EIMS (70 eV) *m/z* (relative intensity) 477 (3, M⁺); [α]_D²⁵ -138° (c 1.05, chloroform).

Example 45**(S)-N'-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(morpholinocarbonyl)benzamide**

4-(Morpholinocarbonyl)benzoic acid (100 mg, 0.43 mmol; described in: *J. Med. Chem.* 1994, 37(26), 4538-4554) and 1,1'-carbonyldiimidazole (76 mg, 0.47 mmol) were

1 dissolved in *N,N*-dimethylformamide (3 mL) and heated to 75 °C for 3.5 h. Additional 1,1'-
2 carbonyldiimidazole (36 mg, 0.22 mmol) was added and the solution was stirred for 30 min.
3 (S)-3-Amino-5-(4-methylpiperazin-1-yl)-3,4-dihydropyran-2H-1-benzopyran (100 mg, 0.40
4 mmol), dissolved in *N,N*-dimethylformamide (2 mL), was added and the reaction mixture
5 was stirred for 18 h at 50 °C. The solvent was evaporated and the residue was partitioned
6 between ethyl acetate (30 mL) and water (5 mL). The organic layer was washed with water
7 (5 mL) and brine (5 mL) and dried (MgSO₄). The solvent was evaporated *in vacuo* giving
8 180 mg of a crude product. Purification by preparative TLC (twice using chloroform/
9 methanol/concentrated ammonia 95:5:0.5 and chloroform/ethanol (saturated with NH₃)
10 12:1 as the eluents afforded 98 mg (53% yield) of the title compound as a white solid: mp
11 222 °C (decomposes); EIMS(70 eV) *m/z* (relative intensity) 464 (68, M⁺); [α]_D²² -12° (c
12 0.44, chloroform).

Example 46

13 (S)-*N*-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-4-(*N,N*-
14 dimethylamino)benzamide
15 To 4-(*N,N*-dimethylamino)benzoic acid (110 mg, 0.56 mmol; described in: U.S.
16 Patent 3,607,918, 1971) was dropwise added thionyl chloride (500 μL, 6.9 mmol). The
17 reaction mixture was stirred at ambient temperature for 1 min and then concentrated *in*
18 *vacuo*. The excess of thionyl chloride was co-evaporated with toluene *in vacuo*. The crude
19 acid chloride was dissolved in methylene chloride (8 mL) and dropwise added to a solution
20 of (S)-3-amino-5-(4-methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran (130 mg, 0.53
21 mmol) and triethylamine (110 μL, 0.80 mmol) in methylene chloride (5 mL) at 0 °C. The
22 reaction mixture was stirred at 0 °C for 30 min and at room temperature for an additional
23 30 min. The solvent was evaporated *in vacuo* giving 300 mg of a crude product.
24 Purification by preparative TLC on silica using chloroform/ethanol (saturated with
25 ammonia) 10:1 as the eluent afforded 120 mg (54% yield) of the title compound as a white
26 solid: mp 219 °C (dec); EIMS (70 eV) *m/z* (relative intensity) 422 (47, M⁺); [α]_D²² -12° (c
27 0.42, chloroform)

Example 47

1 (S)-*N*-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-4-(4-(2-
2 benzoyloxyethyl)-piperazin-1-yl)benzamide
3 (S)-*N*-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-4-(piperazin-1-
4 yl)benzamide (500 mg, 1.2 mmol) was dissolved in *N,N*-dimethylformamide (5 mL) and
5 potassium carbonate (170 mg, 1.3 mmol) was added. To the mixture was added a solution
6 of 2-benzoyloxyethyl mesylate (290 mg, 1.3 mmol) (described in: Beard, C.; Edwards, J.;
7 Fried, J. U.S Patent 3 929 824, 1972) in *N,N*-dimethylformamide (5 mL). The reaction
8 mixture was stirred at 40 °C for 24 h. The solvent was evaporated *in vacuo* giving 950 mg
9 of a crude product. Purification by column chromatography on silica gel using
10 chloroform/methanol/concentrated ammonia 95:5:0.5 as the eluent afforded 154 mg (24%
11 yield) of the title compound as an oil: EIMS (70 eV) *m/z* (relative intensity) 569 (3, M⁺).

Example 48

12 (S)-*N*-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-4-(4-(2-
13 hydroxyethyl)-piperazin-1-yl)benzamide
14 (S)-*N*-(5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl)-4-(4-(2-
15 benzoyloxyethyl)-piperazin-1-yl)benzamide (150 mg, 0.27 mmol) was dissolved in acetic
16 acid (10 mL) and palladium (10%) on carbon (12 mg) was added. Hydrogenation at room
17 temperature and at atmospheric pressure for 14 h followed by filtration and evaporation of
18 the solvent *in vacuo* gave 180 mg of a crude product. The residue was partitioned between
19 methylene chloride (60 mL) and 2 M NH₃ (5 mL) and washed with brine (5 mL). Drying
20 (MgSO₄) the solution and evaporation of the solvent *in vacuo* gave 120 mg of crude
21 material. Purification by preparative TLC on silica using
22 chloroform/methanol/concentrated ammonia 95:5:0.5 as the eluent afforded 37 mg (29%
23 yield) of the title compound as a white solid: mp 211-212 °C; EIMS (70 eV) *m/z* (relative
24 intensity) 479 (8, M⁺); [α]_D²² -26° (c 0.26, chloroform).

PHARMACOLOGY

Electrical field stimulation of [3 H]-5-HT release from occipital cortex of guinea pigs
 [3 H]-5-HT is released by electrical field stimulation from slices of occipital cortex of guinea pigs which have been pre-incubated with [3 H]-5-HT. This release is similar to that caused by nerve stimulation, i.e. exocytotic release from serotonergic nerve terminals, depending on the presence of Ca^{2+} in the incubation medium. The 5-HT release is regulated at the level of the nerve terminals by autoreceptors, in the guinea pigs (like in humans) belonging to the h5-HT_{1B} receptor subtype. Thus, agonists of h5-HT_{1B} receptors reduce the amount of [3 H]-5-HT released by electrical field stimulation whereas the release is increased by antagonists of this receptor type. Testing compounds with this method is accordingly a convenient screening technique for determining the potency and functional effect of new h5-HT_{1B} receptor agonists and antagonists.

Methods and Materials

Buffer composition (mM) NaHCO_3 (25), NaH_2PO_4 , H_2O (12), NaCl (117), KCl (6), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (1.2), CaCl_2 (1.3), EDTA Na_2 (0.03). The buffer is gassed for at least 30 min before use. The pH of the buffer is about 7.2 in the room temperature but it rises to about 7.4 at 37 °C.

Preparation of occipital cortical slices

Guinea pigs (200-250 g) were decapitated and the whole brain was removed. The occipital cortex was dissected and cut to slices 0.4x4 mm with McIlwain chopper machine. The white part of the tissue should be removed carefully with a tweezers before slicing. The slices were incubated in 5 ml buffer in the presence of 5 mM pargyline chloride. After incubation with 0.1 mM [3 H]-5-HT for another 30 min the slices were transferred to a test tube and washed three times with same volume buffer. The slices were transferred to the superfusion chambers with a plastic pipette and were washed for 40 min with the buffer in the presence of uptake inhibitor citalopram 2.5 μM with a flow 0.5 ml/min.

Electrical stimulation of 5-HT release

The superfused buffer was collected in 2 ml/fraction. The slices were stimulated by electricity with a train of pulses of frequency 3 Hz, duration 2 ms and current 30 mA for 3 min at the 4th and 13th fractions. The tested drugs were added from the 8th fraction to the end of experiment.


Results

A first electrical (or K^+) stimulation results in a standard amount of [3 H]-5-HT released (S_1). Before the first and the second stimulation the h5-HT_{1B} antagonist is added to the media which results in a dose depending increase of the release (S_2) after the second stimulation. See Fig. 1.

The S_2/S_1 ratio which is the per cent of released [3 H]-5-HT at the second stimulation (S_2) divided by that of the first stimulation (S_1) was used to estimate drug effects on transmitter release.

CLAIMS

1. A compound having the formula (I)



(I)

wherein

X is N or CH;

Y is NR_2CH_2 , CH_2NR_2 , NR_2CO , CONR_2 , NR_2SO_2 or NR_2CONR_2

wherein R_2 is H or $\text{C}_1\text{-C}_6$ alkyl;

R_1 is H, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl;

R_3 is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl or $(\text{CH}_2)_n\text{-aryl}$,

wherein aryl is phenyl or a heteroaromatic ring containing one or two heteroatoms selected from N, O and S and which may be mono- or di-substituted with R_4 and/or

R_5 ;

wherein R_4 is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, halogen, CN, CF_3 , OH, $\text{C}_1\text{-C}_6$ alkoxy, NR_6R_7 , OCF_3 , SO_3CH_3 , SO_3CF_3 , $\text{SO}_2\text{NR}_6\text{R}_7$, phenyl, phenyl- $\text{C}_1\text{-C}_6$ alkyl, phenoxy, $\text{C}_1\text{-C}_6$ alkyphenyl, an optionally substituted

heterocyclic ring containing one or two heteroatoms selected from N, O, S, SO and SO_2 wherein the substituent(s) is(are) selected from $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, phenyl- $\text{C}_1\text{-C}_6$ alkyl, $(\text{CH}_2)_m\text{OR}_9$ wherein m is 2-6 and

R_9 is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl or phenyl- $\text{C}_1\text{-C}_6$ alkyl, and COR_8 , an optionally substituted heteroaromatic ring containing one or two

heteroatoms selected from N, O and S wherein the substituent(s) is(are) selected from $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl and phenyl- $\text{C}_1\text{-C}_6$ alkyl, or COR_8 ;

wherein R_6 is H, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl;

R_7 is H, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl; and

R_8 is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl, CF_3 , NR_6R_7 , phenyl, a

heteroaromatic ring containing one or two heteroatoms selected from N, O and S or a heterocyclic ring containing one or two heteroatoms selected from N, O, S, SO and SO_2 ;

R_3 is H, OH, CF_3 , OCF_3 , halogen, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkoxy;

and n is 0-4;

as (*R*)-enantiomers, (*S*)-enantiomers or a racemate in the form of a free base or a pharmaceutically acceptable salt or solvate thereof.

2. A compound according to claim 1 wherein Y is NR_2CO or CONR_2 .

3. A compound according to any one of claims 1-2 wherein X is N.

4. A compound according to any one of claims 1-3 wherein R_1 is H or $\text{C}_1\text{-C}_6$ alkyl.

5. A compound according to any one of claims 1-4 wherein R_3 is $(\text{CH}_2)_n\text{-aryl}$.

6. A compound according to any one of claims 1-4 wherein R_3 is $(\text{CH}_2)_n\text{-aryl}$ which is substituted with R_4 , which is an optionally substituted heterocyclic or heteroaromatic ring containing one or two heteroatoms selected from N, O and S, or COR_8 .

7. A compound according to any one of claims 5 and 6 wherein n is 0

8. A compound according to claim 6 wherein R_8 is NR_6R_7 or a heterocyclic ring containing two heteroatoms selected from N and O.

9. A compound according to any one of claims 1- 8 wherein X is N and Y is NR₂CO .

10. A compound according to claim 9 wherein X is N, Y is NR₂CO and R₄ is morpholino or COR₈.

11. A compound which is

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-morpholinobenzamide;

10 (S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-piperidinobenzamide;

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-butoxybenzamide;

15 (S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-trifluoromethylbenzamide;

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-N,N-diethylamino benzamide;

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-trifluoromethoxybenzamide;

20 (S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-piperidin-1-yl)benzamide;

(S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(hexahydro-1,4-diazepin-5-on-1-yl)benzamide, or

25 (S)-N-[5-(4-Methylpiperazin-1-yl)-3,4-dihydro-2H-1-benzopyran-3-yl]-4-(4-benzylpiperazin-1-yl)benzamide

in the form of a free base or a pharmaceutically acceptable salt or solvate thereof.

12. A pharmaceutical formulation comprising as active ingredient a therapeutically effective amount of the compound of any one of claims 1-11 as an enantiomer or racemate

in the form of a free base or a pharmaceutically acceptable salt or solvate thereof optionally in association with diluents, excipients or inert carriers.

13. A pharmaceutical formulation according to claim 12 for use in the treatment of 5 5-hydroxytryptamine mediated disorders.

14. A pharmaceutical formulation according to any one of claims 12 or 13 for use in the treatment of mood disorders, anxiety disorders, personality disorders, obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders, pathological aggression, schizophrenia, endocrine disorders, stroke, dyskinesia, Parkinson's disease, thermoregulatory disorders, pain, hypertension, urinary incontinence or vasospasm; or for growth control of tumors.

15. A compound as defined in any of claims 1-11 for use in therapy.

16. A compound as defined in claim 15 for use in the treatment of disorders in the central nervous system.

20 17. A compound as defined in claim 16 for use in the treatment of mood disorders, anxiety disorders, personality disorders, obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders, pathological aggression, schizophrenia, endocrine disorders, stroke, dyskinesia, Parkinson's disease, thermoregulatory disorders, pain or hypertension.

25 18. A compound as defined in claim 15 for use in the treatment of urinary incontinence or vasospasm or for growth control of tumors.

19. A compound as defined in claim 15 for use in the treatment of 5-hydroxytryptamine mediated disorders.

20. A compound as defined in claim 19 for use as a h5-HT_{1B} antagonist.

21. The use of a compound defined in any of claims 1-11 in the manufacture of a medicament for the treatment of disorders in the central nervous system and/or urinary incontinence or vasospasm; or for growth control of tumors.

22. The use according to claim 21 in the manufacture of a medicament for the treatment of mood disorders, anxiety disorders, personality disorders, obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders, pathological aggression, schizophrenia, endocrine disorders, stroke, dyskinesia, Parkinson's disease, thermoregulatory disorders, pain or hypertension.

23. The use of a compound defined in any of claims 1-11 in the manufacture of a medicament for the treatment of 5-hydroxytryptamine mediated disorders

24. The use according to claim 23 wherein the compound according to any one of claims 1-11 is used as a h5-HT_{1B} antagonist.


25. A method for the treatment of disorders in the central nervous system and/or urinary incontinence or vasospasm or for growth control of tumors by administering to a mammal including man in need of such a treatment a therapeutically effective amount of a compound defined in any of claims 1-11.

26. A method according to claim 25 for the treatment of mood disorders, anxiety disorders, personality disorders, obesity, anorexia, bulimia, premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders, pathological aggression, schizophrenia, endocrine disorders, stroke, dyskinesia, Parkinson's disease, thermoregulatory disorders, pain or hypertension.

27. A method for the treatment of 5-hydroxytryptamine mediated disorder by administering to a mammal including man in need of such a treatment a therapeutically effective amount of a compound defined in any of claims 1-11.

28. A method according to claim 27 wherein the compound according to any one of claims 1-11 is used as a h5-HT_{1B} antagonist.


29. A process for the preparation of the compound of formula I according to claim 1 by acylation, in the case when R₁ is C₁-C₆ alkyl or C₃-C₆ cycloalkyl, Y is NR₂CO, R₂ is hydrogen and X and R₃ are as defined in general formula I in claim 1, of a compound of formula A₁.



with an activated carboxylic acid R₃-CO-Lg, where Lg is a leaving group or by using a carboxylic acid R₃-COOH with an activating reagent

A(II)
acylation, in the case when R₁ is hydrogen, Y is NR₂CO, R₂ is hydrogen, R₃ is a protecting group and X and R₃ are as defined in general formula I in claim 1, of a compound of formula B

69




with an activated carboxylic acid $R_3\text{-CO-Lg}$, where Lg is a leaving group or by using a carboxylic acid $R_3\text{-COOH}$ with an activating reagent, followed by the removal of the protecting group R_6 ;

A(III)


debenzylation, in the case when R_1 is $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, X and R_2 are as defined in general formula I above and R_3 below is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl,

$(\text{CH}_2)_m\text{OH}$ wherein m is 2-6 or COR_8 , of a compound of formula Ia, followed by
a) hydrogenation, b) alkylation, c) alkylation and removal of a protecting group or
d) acylation;

**B(I)**

reacting, in the case when R_1 is $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, Y is CONR_2 , X , R_2 and R_3 are as defined in general formula I above, an activated carboxylic acid of a compound of formula C;


70



with an aniline or amine HNR_2R_3 , or

B(II)

reacting, in the case when R_1 is hydrogen, Y is NR_2CO , R_6 is a protecting group and X , R_2 and R_3 are as defined in general formula I above, an activated carboxylic acid of a compound of formula D




with an aniline or amine HNR_2R_3 , followed by removal of the protecting group R_6 .

C

reaction, in the case when R_1 is $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_3\text{-C}_6$ cycloalkyl, Y is NR_2CONR_2 , R_2 is hydrogen and X and R_3 are as defined in general formula I above, a compound of formula A,

71

1 / 1




(A)

(B)

with a suitable azide in the presence of carboxylic acid, R_3COOH .

30. A compound having the formula




wherein

$X = N$ or CH ;

$Z = NH_2$ or $COOH$;

R_1 is H , C_1-C_6 alkyl or C_3-C_6 cycloalkyl.



INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 98/01603

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 311/58, A61K 31/495, A61K 31/535
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,OK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS-ONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5420151 A (EVA M. HAMMARBERG ET AL), 30 May 1995 (30.05.95), see especially example 83 and column 3 - column 6	1-24, 29-30
A	WO 9109853 A1 (AKTIEBOLAGET ASTRA), 11 July 1991 (11.07.91)	1-24, 29-30
A	WO 9012795 A1 (THE UPJOHN COMPANY), 1 November 1990 (01.11.90)	1-24, 29-30
A	WO 9707120 A1 (SMITHKLINE BEECHAM PLC), 27 February 1997 (27.02.97)	1-24, 29-30

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

- * Special categories of cited documents
- *X* document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *Y* document of particular relevance to the claimed invention, cannot be considered novel or cannot be considered to involve an inventive step when the document is taken into account
- *L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another claim or other document
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed
- *Z* document number of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

1 8 -01- 1999

4 JANUARY 1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson

Telephone No. +46 8 722 25 00

Form PCT/ISA210 (second sheet) (July 1993)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 98/01603

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claim Nos.: 25-28

because they relate to subject matter not required to be searched by this Authority, namely:

A method for treatment of the human or animal body by therapy, see rule 39.1

2. ☐ Claim Nos.:

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim Nos.:

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(c).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This international Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos.:4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA210 (continuation of first sheet) (July 1993)

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/12/98
PCT/SE 98/01603

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/12/98
PCT/SE 98/01603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5420151 A	30/05/95	AU 687115 B	19/02/98
		AU 8069794 A	22/09/95
		BR 9407510 A	26/11/95
		CA 2173257 A	04/05/95
		CN 1136811 A	27/11/96
		CZ 9601114 A	11/09/96
		EP 0725779 A	14/08/96
		FI 961809 A	20/05/96
		HR 940890 A	31/12/95
		HU 73060 A	28/03/97
		HU 9601103 D	00/00/00
		IL 111365 D	00/00/00
		JP 9504287 T	28/04/97
		NO 961687 A	03/06/96
		NZ 275176 A	24/11/97
		PL 314086 A	19/08/96
		SK 43596 A	01/10/96
		US 5500425 A	19/03/96
		US 561610 A	01/04/97
		US 5639772 A	17/06/97
		US 5639784 A	17/06/97
		US 5641807 A	24/06/97
		US 5646309 A	08/07/97
		US 5650524 A	22/07/97
		US 5656557 A	12/08/97
		US 5656558 A	12/08/97
		WO 9511891 A	04/05/95
		ZA 9408353 A	02/05/95
		AT 138918 T	15/06/96
		AU 641204 B	16/09/93
		AU 6977091 A	24/07/91
		CA 2047237 A	23/06/91
		CN 1037438 B	18/02/98
		CN 1052669 A	03/07/91
		CY 1991 A	03/09/97
		CZ 283630 B	13/05/98
		CZ 9006578 A	18/02/98
		DE 69027311 D,1	24/10/96
		DK 460169 T	23/09/96
		EG 19749 A	31/01/96
		EP 0460169 A,B	11/12/91
		SE 0460169 T3	

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
ES 2087993 T	01/08/96		
FI 102177 B	00/00/00		
FI 913943 D	00/00/00		
GR 3020497 T	31/10/95		
HK 55897 A	09/05/97		
HR 920851 A	31/10/95		
HU 211860 B	28/12/95		
HU 9500717 A	28/12/95		
IL 96712 A	31/08/95		
JP 4503682 T	02/07/92		
LT 1730 A	25/08/95		
LV 3967 B	27/05/96		
LV 11894 A,B	20/02/95		
MX 23889 A	01/10/93		
NO 180336 B,C	23/12/96		
PL 164592 B	31/08/94		
PL 164608 B	31/08/94		
PT 96304 A	30/09/91		
RU 2092483 C	10/10/97		
SE 8804361 D	00/00/00		
SG 48043 A	17/04/98		
SI 9012429 A	31/12/97		
WO 9109853 A	11/07/91		
AT 139774 T	15/07/96		
AU 667687 B	04/04/96		
BG 2768492 A	03/05/93		
CA 2118708 A	31/05/95		
CN 1072411 A	15/04/93		
CN 1146592 A	26/05/93		
CY 1992 A	09/04/97		
CZ 9400510 A	05/09/97		
DE 69211857 D,1	16/11/94		
DK 607274 T	24/10/96		
EP 0538222 A	04/11/96		
EP 0607274 A,B	21/04/93		
SE 0607274 T3	27/07/94		
EP 0703229 A	27/03/96		
ES 2089571 T	01/10/96		
FI 941616 A	08/04/94		
GR 3020608 T	31/10/96		
HK 55997 A	09/05/97		
HR 920935 A	31/08/95		
HU 68834 A	28/08/95		
HU 9400993 D	00/00/00		
JP 7500320 T	12/01/95		
MX 9205744 A	01/04/93		
NO 941256 A	07/04/94		
NZ 244618 A	28/10/96		
NZ 270595 A	28/10/96		
PL 171013 B	28/02/97		
SE 9102905 D	00/00/00		
SK 36194 A	07/12/94		

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/12/98

International application No.
PCT/SE 98/01603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5420151 A	30/05/95	WO 9307135 A ZA 9207609 A SE 9202000 D	15/04/93 13/04/93 00/00/00
WO 9109853 A1	11/07/91	AT 138918 T AU 641204 B AU 6977091 A CA 2047237 A CN 1037438 B CN 1052669 A CY 1991 A CZ 283630 B CZ 9006578 A DE 69027311 D, T DK 460169 T EP 19749 A EP 0460169 A, B EP 0460169 T3 ES 2087993 T FI 102177 B FI 913943 D GB 3020497 T HK 55897 A HR 920851 A HU 211860 B HU 9500717 A IL 96712 A JP 4503682 T LT 1730 A LT 3967 B LV 10449 A, B LV 11894 A, B MK 23889 A NO 180336 B, C PL 164592 B PL 164608 B PL 164609 B PT 96304 A RU 2092483 C SE 8904361 D SE 48043 A SI 9012429 A US 5420151 A US 5500425 A US 5616610 A US 5639772 A US 5639784 A US 5641887 A US 5646309 A US 5650524 A US 5656657 A US 5656658 A	15/06/96 16/09/93 24/07/91 23/06/91 18/02/98 03/07/91 05/09/97 13/05/98 18/02/98 24/10/96 23/09/96 31/01/96 11/12/91 01/08/96 00/00/00 00/00/00 31/10/96 09/05/97 31/10/95 28/12/95 31/08/95 02/07/92 25/08/95 27/05/96 20/02/95 20/12/97 01/10/93 23/12/96 31/08/94 31/08/94 31/08/94 30/09/91 10/10/97 00/00/00 17/04/98 31/12/97 30/05/95 19/03/96 01/04/97 17/06/97 17/06/97 24/06/97 08/07/97 22/07/97 12/08/97

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/12/98

International application No.
PCT/SE 98/01603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9012795 A1	01/11/90	AU 627027 B AU 5555290 A CA 2047236 A, C DE 69005625 D, T DK 470176 T EP 0470176 A, B SE 0470176 T3 JP 4504721 T US 5306830 A	13/08/92 16/11/90 28/10/90 11/05/94 25/04/94 12/02/92 20/08/92 26/04/94
WO 9707120 A1	27/02/97	GB 9516456 D GB 9606632 D GB 9606633 D	00/00/00 00/00/00 00/00/00